

Sản xuất các axít béo thiết yếu không bão hòa đa chuỗi dài bởi sinh trưởng quang tự dưỡng, hợp dưỡng và dị dưỡng ở các vi sinh vật quang hợp và không quang hợp: tổng quan

Phan Minh Tâm¹, Trần Nguyên Vũ^{2,*}

¹*Khoa Khoa học Sinh học, Trường Đại học Nông Lâm Thành phố Hồ Chí Minh, Việt Nam*

²*Khoa Sinh học – Công nghệ Sinh học, Trường Đại học Khoa học Tự nhiên
Thành phố Hồ Chí Minh, Việt Nam*

*Ngày nhận bài: 25/08/2023; Ngày sửa bài: 05/12/2023;
Ngày nhận đăng: 16/02/2024; Ngày xuất bản: 28/02/2024*

TÓM TẮT

Một số vi sinh vật quang hợp và không quang hợp như vi tảo, sinh vật nguyên sinh hoặc nấm, được biết đến như là nguồn sản xuất tự nhiên của các axit béo không bão hòa đa chuỗi dài (LC-PUFA). Trong đó, một số loài được biết đến như là những sinh vật dị dưỡng bắt buộc, hợp dưỡng hoặc quang tự dưỡng bắt buộc. Tuy nhiên, ngày càng có nhiều loài vi tảo, trước đây được biết là sinh vật quang tự dưỡng bắt buộc, nhưng nay được xác định là sinh vật hợp dưỡng hoặc dị dưỡng. Các con đường sinh tổng hợp và điều kiện nuôi cấy của các vi sinh vật này được so sánh để làm nổi bật các yếu tố ảnh hưởng đến quá trình sản xuất và phân phối LC-PUFA trong tế bào. Sản xuất LC-PUFA đã được cải thiện bằng cách lựa chọn quy trình nuôi cấy và chủng vi sinh vật. Các phân tích về sản lượng chuyển đổi và năng suất của LC-PUFA trong nuôi cấy quang tự dưỡng, hợp dưỡng và dị dưỡng làm sáng tỏ hiệu suất sản xuất LC-PUFA bởi các sinh vật quang hợp và không quang hợp.

Từ khóa: *Vi tảo, axit béo không bão hòa đa (PUFA), sinh trưởng quang tự dưỡng, sinh trưởng hợp dưỡng, sinh trưởng dị dưỡng.*

**Tác giả liên hệ chính.*

Email: tnvu@hcmus.edu.vn

Photo-autotrophic, mixotrophic and heterotrophic production of essential long chain polyunsaturated fatty acids in photosynthetic and non-photosynthetic microorganisms: a review

Tam Minh Phan¹, Nguyen-Vu Tran^{2,*}

¹Faculty of Biological Sciences, Nong Lam University Ho Chi Minh City, Vietnam

²Faculty of Biology - Biotechnology, Ho Chi Minh City University of Science, Vietnam

Received: 25/08/2023; Revised: 05/12/2023;

Accepted: 16/02/2024; Published: 28/02/2024

ABSTRACT

Some photosynthetic and non-photosynthetic microorganisms such as microalgae, stramenopiles or fungi, are known as natural producers of long chain polyunsaturated fatty acids (LC-PUFAs). Among those, some species are known as obligate heterotrophs, mixotrophs or obligate phototrophs. However, more and more microalgal species, previously reported as obligate photo-autotrophs, are now identified as mixotrophs or heterotrophs. The biosynthetic pathways and cultivation conditions of these microorganisms are compared to highlight the factors influencing production and distribution of LC-PUFAs in the cells. LC-PUFA production has been improved by the choice of cultivation processes and microorganism strains. Analyses of the conversion yields and productivities of LC-PUFAs in photo-autotrophic, mixotrophic and heterotrophic cultivation elucidate the performance of LC-PUFA production by photosynthetic and non-photosynthetic organisms.

Keywords: Microalgae, polyunsaturated fatty acid (PUFA), photoautotrophic growth, mixotrophic growth, heterotrophic growth.

Abbreviations: LC-PUFAs: long chain polyunsaturated fatty acids; GLA: gamma-linolenic acid; ARA: arachidonic acid; EPA: eicosapentaenoic acid; DHA: docosahexaenoic acid; H: heterotrophic growth; M: mixotrophic growth; P: photoautotrophic growth; DO: dissolved oxygen; PKS: polyketide synthase; ROS: reactive oxygen species; TAG: triacylglycerol; MAG: monoacylglycerol; PC: phosphatidylcholine; PE: phosphatidylethanolamine; MGDG: monogalactosyldiacylglycerol; DPG: diphosphatidylglycerol; SQ: sulfoquinovosyldiglyceride; mg/L/d: milligram.liter⁻¹.day⁻¹.

1. INTRODUCTION

During last decades, there was growing interest in supplying unsaturated lipids to animal and human.¹ Indeed, some particular unsaturated lipids have been shown to benefit to animal and human health.²⁻⁴ These lipids are provided either by food intake or by extracts e.g. fish oils, olive oils, soybean oils, canola oils,

flaxseed oils.⁵ These sources are dependent on seasonal variations or the availability of natural resources. Moreover, fish oils are dependent on risks of contamination by xenobiotics^{6,7} and of unpleasant smell and taste. Therefore, these drawbacks were the reason why relatively recent bioprocesses for production by microorganisms in bioreactors have been developed.^{8,9} One of

*Corresponding author.

Email: tnvu@hcmus.edu.vn

the first industrial applications was the DHA production by a microalga *Cryptocodinium cohnii*.¹⁰ Other photosynthetic and non-photosynthetic microorganisms were also found to produce unsaturated lipids.^{11,12}

The unsaturated lipids having 18 carbons or more in length with two or more double bonds were characterized as LC-PUFAs. Gamma-linolenic acid (GLA), arachidonic acid (ARA), eicosahexaenoic acid (EPA) and docosahexaenoic acid (DHA) are the essential LC-PUFAs which have been found in different sources as plant, fish, egg... In recent years, these essential LC-PUFAs have been produced by microorganisms.¹ These microorganisms include some microalgae, lower fungi, and bacteria which were grown in photoautotrophic, mixotrophic or heterotrophic condition (aerobic or anaerobic condition). These conditions also affect LC-PUFA content in the cells.¹³ LC-PUFAs produced by microorganisms were naturally esterified as glycolipids, phospholipids, and neutral lipids which were constituted of membrane compositions.

Oxygen plays an important role for most life on the earth because all higher organisms are aerobioses. Oxygen is indispensable for aerobioses. In cells, oxygen can transform into more reactive forms, Reactive Oxygen Species (ROS) which are toxic to cells. Under oxidative stresses, cells have to maintain the balance of the production between ROS and antioxidant enzymes. However, when the generation of ROS overtakes antioxidant enzymes, the damage of lipids, proteins and nucleic acids occurs.¹⁴ Besides the effects of cell component damage, oxygen also participates in biosynthesis and metabolism of cell compositions. During biosynthesis of LC-PUFAs, oxygen plays a role as an electron acceptor in reduction of fatty acids to form the double bonds.

In cultivation, some factors were useful for overproduction of LC-PUFAs such as temperature, pH, salinity, light,^{15,16} oxygen tension.¹⁷ Besides the environmental factors, nutritional factors also affect LC-PUFA

biosynthesis. Organisms used in culture can be photoautotrophs, mixotrophs or heterotrophs. Photoautotrophs are organisms that obtain energy from light and carbon source from CO₂ to synthesize organic compounds in the cells while mixotrophs can use energy from light and carbon sources from organic compounds.¹⁸ Heterotrophs obtain energy and carbon sources from organic compounds.

For improvement of LC-PUFA production, many different types of bioreactors were used from pilot to industrial scales.^{9,19,20} Depending on the value of the desired products, photosynthetic production can be carried out in open systems and closed systems (photobioreactor).¹⁶ For heterotrophic culture, the classical enclosed bioreactors were used. Industrial reactors and data are available for some photoautotrophic and heterotrophic processes but a lesser point for mixotrophic process.

The objective of this review is to compare the performance of LC-PUFA production by photosynthetic and non-photosynthetic microorganisms.

2. MICROBIAL SOURCES OF LC-PUFA

LC-PUFAs are produced by various microorganisms, from prokaryotes to eukaryotes. They could be also classified either as non-photosynthetic microorganisms or photosynthetic microorganisms (Table 1).

Non-photosynthetic LC-PUFA producers constitute a large group of microorganisms that obtain energy and carbon from organic compounds. This group includes bacteria, fungi, fungus-like microorganisms and some microalgae. Some bacteria have been shown to produce EPA²¹ or DHA.²² These bacteria are *Shewanella*²²⁻²⁵ and *Moritella*.²⁶ Some fungi can produce GLA, ARA or EPA. GLA-producing fungi include genera *Mortierella*,²⁷⁻²⁹ *Cunninghamella*,^{1,30-32} *Pythium*³³ and *Mucor*.^{1,34,35} ARA-producing fungi include genera *Mortierella*^{36,37} and *Pythium*.³⁸ These fungal genera are also able to produce EPA.³⁹⁻⁴¹ Other groups include fungus-like microorganisms,

known as Stramenopiles, characterized as particularly marine stramenopilan protists belonging to the class Labyrinthulomycetes. They differ from fungi in composition of their cell walls (absence of chitin) and rhizoids.⁴² Stramenopiles produce high levels of DHA up to 65.9% of total fatty acids.⁴³ The genera *Thraustochytrium*,⁴⁴⁻⁴⁷ *Schizochytrium*⁴⁸⁻⁵⁰ and *Aurantiochytrium*^{51,52} are representatives for DHA production. Finally, the microalga *Cryptocodonium* is included in the non-photosynthetic microorganisms, as it is an obligate heterotroph. This microalga can produce high amounts of DHA up to 63.2% of total fatty acids.⁵³

Photosynthetic microorganisms are the microorganisms for which energy is provided by light and carbon either by inorganic carbon (photoautotrophy) or by organic carbon sources (mixotrophy). This group includes cyanobacteria and microalgae. Some

of them can grow mixotrophically and/or heterotrophically.⁵⁴⁻⁵⁷ Cyanobacterium *Spirulina* only biosynthesizes GLA while some other microalgae can biosynthesize ARA, EPA or DHA. Not all cyanobacteria or microalgae species can produce LC-PUFAs, only certain cyanobacteria or microalgae can biosynthesize these LC-PUFAs. Microalgae can produce ARA such as *Porphyridium*,⁵⁸ *Parietochloris*,⁵⁹⁻⁶¹ *Euglena*⁶² and *Galdieria*.⁶³ Microalgae produce EPA such as *Monodus*,⁶⁴⁻⁶⁶ *Porphyridium*,^{58,67} *Phaeodactylum*,^{68,69} *Nannochloropsis*,⁷⁰⁻⁷² *Navicula*,^{73,74} *Nitzschia*,⁷⁵⁻⁷⁸ *Skeletonema*.⁷⁹ Some microalgae such as *Rhodomonas*,⁸⁰ *Isochrysis*⁸¹ or *Pavlova*⁸² contain both EPA and DHA in their cells. *Porphyridium* produces both ARA and EPA.⁵⁸ *Nannochloropsis*,⁷² *Nitzschia laevis*¹³ can produce EPA in 3 modes of nutrition. *Spirulina* can also grow and produce GLA in 3 modes of nutrition.^{57,83}

Table 1. Distribution of LC-PUFAs in microorganisms.

Microorganisms	Strains (*)	GLA	ARA	EPA	DHA
Non-photosynthetic	Bacteria	<i>Shewanella oneidensis</i> ATCC 700550 ²³		+	
		<i>Shewanella putrefaciens</i> MAC1 ²⁴		+	
		<i>Shewanella baltica</i> ²⁵		+	
		<i>Shewanella morphae</i> ²⁵		+	
		<i>Moritella marina</i> ATCC 15381 ²⁶			+
	Fungi	<i>Mucor circinelloides</i> ¹	+		
		<i>Mucor mucedo</i> ¹	+		
		<i>Mucor rouxii</i> ³⁴	+		
		<i>Mucor inaqisporus</i> ³⁵	+		
		<i>Cunninghamella elegans</i> CCF1318 ¹	+		
		<i>Cunninghamella echinulata</i> CCRC 31840 ³⁰	+		
		<i>Pythium debaryanam</i> ³³	+		
		<i>Pythium ultimum</i> ³⁸		+	+
		<i>Pythium irregularare</i> ⁸⁴			+
		<i>Mortierella elongata</i> NRRL 5513 ⁴¹			+
		<i>Mortierella isabellina</i> ²⁹	+		
		<i>Mortierella ramanniana</i> MM15-1 ²⁸	+		
		<i>Mortierella alpina</i> ATCC 16266 ³⁷		+	
		<i>M. alpina</i> ATCC 32222 ⁴⁰			+

	Fungus-like protists	<i>Thraustochytrium aureum</i> ATCC 34304 ⁴⁴				+
		<i>Thraustochytrium roseum</i> ATCC 28210 ⁴⁷				+
		<i>Schizochytrium mangrovei</i> FB3 ⁸⁵				+
		<i>Schizochytrium limacinum</i> SR21 ⁴⁹				+
		<i>Aurantiochytrium mangrovei</i> MP2 ⁵¹				+
		<i>Aurantiochytrium limacinum</i> mh0186 ⁵²				+
Photosynthetic	Microalgae	<i>C. cohnii</i> ATCC 30772 ⁸⁶⁻⁸⁸				+
		<i>Spirulina platensis</i> UTEX 1928 ⁵⁷	+			
	Microalgae	<i>Porphyridium cruentum</i> SAG 1380-1a ⁵⁸		+	+	
		<i>Parietochloris incisa</i> ⁵⁹⁻⁶¹		+		
		<i>Skeletonema costatum</i> ⁷⁹			+	
		<i>Monodus subterraneus</i> Petersen UTEX 151 ⁶⁶			+	
		<i>Nitzschia laevis</i> ⁷⁵			+	
		<i>Nannochloropsis</i> sp. ⁷²			+	
		<i>Navicula saprophila</i> ^{73,74}			+	
		<i>Phaeodactylum tricornutum</i> UTEX 640 ⁶⁸			+	
		<i>Glaucoctysis nostochinearum</i> ⁶²			+	
		<i>Cyanophora paradoxa</i> ⁶²			+	
		<i>Euglena gracilis</i> ⁶²		+	+	

(*) Representative strains associated with applied biotechnological studies.

3. OXYGEN AND LC-PUFABIOSYNTHESIS

There are two different biosynthetic pathways of LC-PUFAs in microorganisms: anaerobic and aerobic pathways (Figure 1).⁸⁹ This classification is based on the oxygen dependence of PUFA biosynthesis reactions.

The term “anaerobic pathway” does not mean that the pathway only occurs in anaerobic condition. It can operate in the presence of oxygen but oxygen was not used for formation of double bonds. The formation of double bonds in this pathway is carried out by a dehydration to

remove a water molecule from hydroxyacyl-acyl carrier protein (ACP).

The anaerobic pathway was found in some non-photosynthetic microorganisms such as some bacteria and fungus-like microorganisms. LC-PUFA biosynthesis in some bacteria is generally carried out by PKS-like system.⁹⁰ To date, intermediates of this pathway have not yet been determined in detail. Some fungus-like microorganisms such as *Thraustochytrium* and *Schizochytrium* carried out two pathways for LC-PUFA biosynthesis: Polyketide Synthase (PKS) anaerobic pathway and aerobic pathway.⁹¹ Results of studies in *C. cohnii* indicated that DHA biosynthesis by this microalga was not catalyzed by the desaturases and elongases in aerobic pathway⁹² but would be made by anaerobic PKS pathway.⁹³

In aerobic pathway, molecular oxygen is required in biosynthesis of unsaturated fatty acids. It acts as an electron acceptor for double bond formation in the unsaturated fatty acids. This process is catalyzed by the desaturase enzymes which remove two H atoms from saturated or unsaturated fatty acids to form double bonds in these fatty acids. These two H atoms combine with the O atom of O₂ to form H₂O. The other O atom is reduced by cytochrome b₅.⁹⁴⁻⁹⁶

The aerobic pathway occurs in microorganisms such as cyanobacteria, fungi and microalgae. Desaturases use molecular oxygen to form double bonds for unsaturated fatty acids production. This biosynthetic pathway has been studied in photosynthetic microorganisms by using ¹⁴C labelled intermediates. In the red microalga *P. cruentum*, it was shown that externally supplied unsaturated fatty acids were assimilated in mixotrophic cultures. Two routes of EPA biosynthesis from linoleic acid (C18:2,

n-6) precursors have been suggested: one route along n-6 pathway and another along n-3 pathway.⁹⁷ In diatom *P. tricornutum*, four routes of EPA biosynthesis were revealed by use the radiolabeled intermediates. Two routes pass along n-3 pathway, one route pass along n-6 pathway and other route pass along both n-3 and n-6 fatty acids as intermediates.⁹⁸ In *Parietochloris incisa*, ARA biosynthesis is carried out in cytoplasm and in chloroplast.⁹⁹ In *Pavlova* and *Isochrysis*, DHA was synthesized by aerobic pathway. Gene *IgASE1* which encoded an elongating enzyme in aerobic pathway was identified in *I. galbana* and expressed in yeast.¹⁰⁰ In addition, the genes *pavELO* (*Pavlova*), which catalysed conversion of EPA into DPA and *IgD4* (*Isochrysis*) catalysed conversion of DPA into DHA, were also identified and transformed into yeast cells. The yeast cells were cultured with exogenously supplied EPA and they can synthesize DHA from EPA.¹⁰¹ Many desaturase and elongase genes in *M. alpina* have been isolated and characterized.¹⁰² This indicated that PUFA biosynthesis in this fungus is carried out by aerobic pathway. Other proof showed that when using SAN 9785, the inhibitor of desaturases, GLA content in the cells of *Spirulina* was significantly influenced.⁶⁷

Using intermediates of the biosynthetic pathways seems to increase the content of LC-PUFAs. The ARA content of *Porphyridium* was 23.2% of total fatty acids after addition of linoleate to the culture medium compared with 16.8% in the control.⁹⁷ Similarly, GLA content was 36.0% of total fatty acids when *Spirulina* was grown with linoleate, against 20.4% in the control.¹⁰³ In the culture of *Mortierella* for EPA production, linseed oil contains α -linolenic acid (precursor for EPA biosynthesis) as a major fatty acid (58%) was supplemented in the medium, resulted in an increase of EPA content.⁴⁰




Figure 1. Pathway of LC-PUFA biosynthesis.

4. FACTORS AFFECTING LC-PUFA PRODUCTION

4.1. Nutritional factors

4.1.1. Carbon sources

Carbon constitutes from 49 to 57% of biomass and about 80% of the LC-PUFAs, provided either by inorganic forms such as CO_2 or HCO_3^- for carbon skeletons in photoautotrophic growth or by organic forms as energy sources and carbon skeletons in mixotrophic or heterotrophic growth.

Glucose is the most commonly used carbon source for heterotrophic growth of microorganisms. However, the suitable carbon sources for growth and LC-PUFA production

would be different among microorganisms. Starch and maltose were the suitable carbon sources for DHA production by *T. aureum* ATCC 34304.⁴⁴ Starch was also the preferred carbon source for GLA production by *C. echinulata* CCRC 31840³⁰ or for ARA production by *M. alpina* ATCC 32222,²⁸ whereas glucose was optimal for DHA production by *Thraustochytrium* sp. KK17-3.¹⁰⁴ Glucose was also the suitable carbon source for growth and EPA production by *P. irregulare* ATCC 10951³⁹ or for GLA production by *Mucor hiemalis* M4.¹⁰⁵ Glucose and starch were suitable carbon sources for ARA production by *Mortierella alliacea* YN-15.¹⁰⁶ Although ARA yield was highest with glycerol, ARA content in lipids was quite low. Thus, glucose was the best choice for ARA production by *M. alpina* ATCC

16266.³⁷ In *M. alpina* CBS 528.72, glucose gave the optimal growth and total lipid content but rhamnose gave a higher ARA content in total fatty acids.¹⁰⁷ The complexe sources as rice bran, wheat bran, peanut meal, sweet potato, linseed oil, soybean oil,... were also investigated for ARA production by *M. alpina* ATCC 32222.²⁸ Marine microalga *C. cohnii* could utilize acetic acid,¹⁰⁸ ethanol,⁸⁷ carob pulp¹⁰⁹ for DHA production.

Some microalgae as *N. laevis* UTEX 2047 could mixotrophically grow with glucose¹³ or *Nannochloropsis* sp. could utilize glucose or ethanol⁷² for EPA production. *P. tricornutum* UTEX-640 could also grow and produce EPA in mixotrophic conditions with various carbon sources. Glycerol was found as the most suitable carbon source for growth and EPA production by this microalga.^{55,56} *Navicula saprophila* could grow with acetate in mixotrophic condition.^{73,74}

In the photoautotrophic culture of *P. tricornutum* UTEX 640, the different concentrations of CO₂ were examined for growth and EPA production. The optimal biomass and EPA yield were 2.5 g/L and 87.5 mg/L, respectively, obtained at 1% CO₂.⁶⁸ In the 4-day cultivation of *Nannochloropsis* sp., 2% CO₂ was supplied 12 h prior to the end of the exponential growth gave the highest EPA yield and productivity which were 340 µg/L and 126 µg/L/d, respectively. This productivity was twice as high as that in ambient air.¹¹⁰ The elevation of CO₂ concentration (350 to 2800 µL/L CO₂) in photoautotrophic culture resulted in an increase of EPA content (21.9% to 25.3% of total fatty acids) in *Nannochloropsis* sp.¹¹¹ *M. subterraneus* UTEX 151 was cultured at two different concentrations of CO₂ (1% and 5%). EPA content in total fatty acids obtained at 1% CO₂ was higher than that obtained at 5% CO₂.⁶⁴

4.1.2. Nitrogen sources - nitrogen starvation

Nitrogen constitutes from 8 to 12% of biomass. LC-PUFA producing microorganisms can grow on organic or inorganic nitrogen sources. *T.*

aureum ATCC 34304 could utilize organic nitrogen sources as tryptone, peptone, malt extract, yeast extract and sodium glutamate. Cells were grown with yeast extract gave 5.0 g/L biomass and 247.7 mg/L DHA yield while those were 3.8 g/L (biomass) and 269.6 mg/L (highest DHA yield) with sodium glutamate.⁴⁴ Tryptone was the most suitable for DHA production by *Thraustochytrium* sp. KK17-3 with 232.8 mg/L DHA.¹⁰⁴ *M. alpina* LPM 301 was grown in the medium with urea or potassium nitrate as nitrogen sources for ARA production. ARA yield was 4.5 g/L after 189 h of cultivation with potassium nitrate and 4.2 g/L after 210 h with urea.¹¹² Some other strains of *Mortierella* fungi have been studied with the inorganic and organic nitrogen sources. Yeast extract was found as the best nitrogen source for growth and ARA production by *M. alpina*.¹¹³ Furthermore, combination of soluble starch 120% and the mixture (2:1, wt/wt) of KNO₃ and yeast extract were the best nitrogen sources for ARA production by *M. alpina* ATCC 32222.²⁸ Ammonium hydroxide was used in the culture of *M. alpina* DSA-12 as the nitrogen source and pH control.¹¹⁴ Ammonium nitrate was found as suitable nitrogen source for GLA production by *C. echinulata*.³⁰ *N. laevis* UTEX 2047 was heterotrophically cultivated with glucose. Nitrate, ammonium and urea were investigated for growth and EPA production. Biomass and EPA yield were over 4 g/L and 90 mg/L, respectively, obtained with nitrate or urea but only 1.24 g/L biomass and 21.58 mg/L EPA with ammonium.⁷⁶ Combination by the ratio 32:1 of glucose and mixture (1:2.6:1.3) of nitrate, tryptone and yeast extract was optimal for EPA production by *N. laevis* UTEX 2047.⁷⁸

Mixotrophic cultivation was carried out with *P. tricornutum* UTEX 640 in the presence of glycerol with urea or nitrate. The best results were 1.52 g/L/d biomass and 43.13 mg/L/d EPA obtained with 0.01 M urea in fed-batch.⁵⁶

In the photoautotrophic cultivation of *P. tricornutum* UTEX 640, urea was the optimal nitrogen source for EPA production.⁶⁸ Nitrate,

nitrite and urea were utilized in the culture of *I. galbana*. DHA content in total fatty acids was highest (14.13%) obtained with urea at early stationary phase.¹¹⁵ $(\text{NH}_4)_2\text{HPO}_4$ was the suitable source of nitrogen for growth and GLA production by *S. platensis*.¹¹⁶

I. galbana CCAP 927/1 was cultivated in nitrate starvation, DHA content in total fatty acids increased from 1.19 to 4.52% from 2nd day to 5th day of cultivation.¹¹⁷ Nitrogen starvation induced an increase in ARA content over 60% of total fatty acids in *P. incisa*.⁶⁰

4.1.3. Phosphorus sources - phosphorus starvation

Phosphorus participates in the energy transfer within cells and constitutes about 5-6% of phospholipids.

In fungus *P. irregulare* ATCC 10951, optimal EPA production (about 31 mg/L) was obtained at 0-3 mM phosphate. The increase of phosphate concentrations (6 – 24 mM) resulted in a decrease of EPA yield.³⁹

Effects of phosphate (0.05-0.5 g/L) were also examined on the growth and EPA production by *P. tricornutum* UTEX 640. Little change in biomass was observed in this range of phosphate concentrations but EPA yield was higher at phosphate levels of 0.1 – 0.5 g/L.⁶⁸

Phosphorus starvation was studied by 7-day cultivation of *P. tricornutum* in the phosphorus-deficient medium (no phosphate was added). A comparative control was made in parallel with 6.9 mg/L NaH_2PO_4 . Results indicated that EPA content in total fatty acids decreased from 26.8% to 6.9% in the condition of phosphorus deficiency.¹¹⁸ Other study showed that EPA content decreased from 28.2 to 15.5% mol of fatty acids when decreasing phosphate concentration (K_2HPO_4) from 175 to 0 μM in the 4-day cultivation of *M. subterraneus*.¹¹⁹

4.1.4. Silicate

Silicate is an essential nutrient for diatom growth because cells need silicate to form their frustules.

N. laevis UTEX 2047 was heterotrophically grown with glucose and silicate. The highest EPA yield (131 mg/L) was obtained at 20 g/L glucose and 32 mg/L silicate while the highest EPA productivity was 15.1 mg/L/d at 20 g/L glucose and 64 mg/L silicate.⁷⁵

In photoautotrophic conditions, the range of silicate from 8.8 – 176 μM has been examined for EPA production in the culture of *Nitzschia inconspicua*. Results showed that there was not significantly change in EPA content (about 4.0% of total fatty acids) and EPA yield (about 0.2 mg/L).¹²⁰ Similarly, the photoautotrophic growth of *P. tricornutum* was not significantly different in the levels of 0 to 50 mg/L silicate. Increase of silicate levels from 50 to 500 mg/L resulted in reducing growth (2.6 to 1.8 g/L biomass) and EPA content (72.5 to 35.0 mg/L EPA).⁶⁸

4.2. Environmental factors

4.2.1. Temperature

Optimal temperature for growth is often different from optimal temperature for LC-PUFA accumulation. The increase of LC-PUFA contents at low temperature is attributed to the cells maintaining fluidity of membranes by biosynthesizing more LC-PUFAs.

Effect of temperature on production of ARA and EPA was studied in *P. ultimum*. The optimal temperature for ARA and EPA production was 25 °C.³⁸ This temperature was also found as the most suitable temperature for ARA accumulation in *M. alpina*¹⁰⁷ and *T. roseum* ATCC 28210 for DHA production.⁴⁵ Highest DHA content in total fatty acids was found when *S. limacinum* OUC88 was cultured at 16 – 23 °C.¹²¹ In *Aurantiochytrium* sp. strain mh0186, cells grew well at 15 – 30 °C, but weakly at 10 °C. The amount of DHA in total fatty acids was highest at 10 °C. The DHA yield was similar at 15 – 30 °C and was significantly higher than those at 10 and 35 °C.¹²² Similarly, *Shewanella* was cultivated at 10, 15 and 25 °C. The cells accumulate with the highest concentration of EPA (6.3% of total lipids) at 10 °C. At 25 °C,

EPA concentration in dry weight is lower (1.5% of total lipids).²⁵ *Galdieria* cells accumulate higher concentrations of PUFAs at 25 °C when compare to 45 °C.⁶³

A range of temperature from 10 to 30 °C was investigated in the culture of *P. tricornutum* 2038. Growth was inhibited at 30 °C, slow at 25 °C and optimal at 20 °C. EPA content in dry weight was highest at 10 °C.¹²³ However, optimal temperature for biomass and EPA production by *P. tricornutum* UTEX 640 was found at 21.5 – 23 °C in the study of Yongmanitchai and Ward (1991).⁶⁸ The effect of temperature on GLA content was also studied in *S. platensis* UTEX 1928. The suitable temperature for GLA accumulation was from 25 to 33 °C.¹²⁴ The optimal EPA production was obtained at 8 °C in the culture of *Porphyridium purpureum* 1380-1b.¹²⁵

4.2.2. pH

Generally, heterotrophic cultures were related to acid pH conditions. pH in the range of 5.5 – 6.5 was suitable for biomass and ARA production by *M. alpina*. Maximal ARA content in total fatty acids was obtained at initial pH 6.5.¹⁰⁷ ARA yield was highest at initial pH 6.0.¹²⁶ Initial pH 6 was also favourable for DHA production by *Thraustochytrium*.^{44,45} Optimal growth and EPA production were obtained at initial pH 6 – 7 in culture of *P. irregulare*.³⁹ The highest DHA content in total fatty acids was 56.8% at initial pH 7.2 in *C. cohnii* ATCC 30556.¹²⁷

In *R. salina*, the concentrations of EPA and DHA accumulated in dry weight when cultivated at pH 8.5 are 0.8% and 0.3%, respectively, compare to 0.6% and 0.2% at pH 7.⁸⁰

Yongmanitchai and Ward have found the maximal EPA yield (93.1 mg/L) at initial pH 7.6 in the photoautotrophic culture of *P. tricornutum* UTEX 640.⁶⁸ The range of pH from 5.0 to 8.5 was tested for EPA production in *P. purpureum* 1380-1b. The highest EPA yield (1.79 mg/L) was obtained at pH 7.6.¹²⁵

4.2.3. Salinity

Some studies relating to effects of salinity on growth and LC-PUFA production have been investigated. A wide tolerance to salinity was found in *S. limacinum* when this fungus-like microorganism was cultured in the salinity range from 0% to 200% that of seawater. In the optimal range of salinity for growth (50 – 200% of seawater), there was little change in dry cell weight. Although this strain could grow at 0% of salinity, the growth was lower than those at the optimal range of salinity.⁴⁹ *Thraustochytrium* sp. showed a slight resistance to high salinity, up to 200% that of seawater. The optimal salinity for growth and DHA production was 75% that of seawater.¹⁰⁴ Sea salt from 2 – 50 g/L was also examined for growth and DHA production by *Thraustochytrium* sp. The highest biomass (24.7 g/L) and DHA yield (4.6 g/L) were obtained at 2 g/L NaCl.⁴⁶ The optimal concentration of NaCl for DHA production by *C. cohnii* ATCC 30556 was 9 g/L.¹²⁸

In *P. lutheri* SMBA60, NaCl concentrations from 5 to 45 g/L were examined for EPA and DHA production. The highest EPA (about 4.7 mg/L) and DHA yield (about 2.6 mg/L) were obtained at 5 – 15 g/L NaCl.¹²⁹ *P. tricornutum* UTEX 640 gave the highest EPA yield at 0 – 10 g/L NaCl.⁶⁸ *Spirulina* was cultivated in the range 0 – 3.5 g/L NaCl, GLA content increased as NaCl level was raised to 0.6 g/L and then it decreased. GLA yield was highest (27 µg/mL) at 0.2 g/L NaCl concentration.¹³⁰

4.2.4. Light

Light also stimulates the growth and DHA production in *T. aureum*. Biomass and DHA yield in light exposed cultures were 70.4 g/L and 269.6 mg/L, respectively, higher than those in dark cultures.⁴⁴ After that, some cultivations of *Thraustochytrium* for DHA production were carried out under light by other authors.^{45,131}

Light affects growth and fatty acid composition of microorganisms, especially the photosynthetic ones. *P. lutheri* SMBA 60 was

grown in semi-continuous cultures at the different light intensities 9, 19 and 30 W/m². The highest EPA and DHA productivities were obtained at 19 W/m².¹³² Effects of intensities and photoperiods on fatty acid production by *I. galbana* have also been studied.¹³³ Percentage of EPA in total fatty acids and in dry weight were 35.7% and 4.4%, respectively when *M. subterraneus* was grown at 90 µmol photon/m²/s, which was higher than those at 170 µmol photon/m²/s.⁶⁴ GLA content in total fatty acids of *S. platensis* increased from 31.1 to 36.0% when increased the light intensity from 860 to 1400 µmol photon/m²/s.¹²⁴

4.2.5. Culture age

Effect of culture age on ARA production by *M. alpina* I₄₉-N₁₈ was investigated. ARA yield increased and was maximal at the 6th day, and then decreased.¹¹³ The GLA yield was also maximal after 5 – 6 days in the culture of *C. echinulata* CCRC 31840.³⁰

In photoautotrophic culture of *Pavlova viridis*, EPA and DHA content in late exponential phase (4 days) were 22.1 and 3.5 mg/g biomass, respectively, and decreased in linear phase (7 days) and stationary phase (13 days).¹³⁴

4.2.6. Dissolved oxygen

Oxygen constitutes from 27 to 32% of biomass and about 10% of the LC-PUFAs. The levels of DO affected growth and LC-PUFA production in heterotrophic culture of various microorganisms. *C. cohnii* gave higher DHA yield when cultured at DO of 10 – 50% of air saturation level.¹⁷ In *S. limacinum* SR21, the culture was carried out in two stages, the first stage for biomass production where concentration of dissolved oxygen at 50% whereas at 10% for DHA production in the second stage.⁴⁹ DHA content of total fatty acids was 30.6% and 40% at 40% DO and 5% DO, respectively.¹³⁵ *Mucor rouxii* ATCC 24905 was shifted from anaerobic to aerobic conditions resulted in an increase of biomass and fatty acid content. Oxygen induced the expression of Δ⁹-, Δ¹²- and Δ⁶- desaturase genes resulted in an increase of unsaturated fatty acids.¹³⁶

5. DISTRIBUTION OF LC-PUFAS IN LIPID CLASSES

Distribution of LC-PUFAs in lipid classes is various among microorganisms. The nutritional and environmental factors affect the distribution of LC-PUFAs in the cells. Information on LC-PUFA localization in the lipid classes is determinant for the purification process.

In *C. echinulata* ATHUM 4411, GLA distribution depended on developmental stages. GLA content in PC remained over 20% of total fatty acids in mid exponential, late exponential and stationary phase whereas that was changed in other lipid classes. ARA content in dry weight increased in non-polar lipids but decreased in polar lipids through growth phases.³² The distribution of LC-PUFAs in lipid classes in *M. alpina* SC9 was influenced by salinity. TAG was the dominant lipid class of the cells (261.16 mg/g) which contained the highest proportion of ARA (30.29% of total fatty acids). When the cells were cultured at 20 g/L NaCl, TAG content increased 296.55 mg/g but ARA content decreased 21.24%.¹³⁷ In *N. laevis* UTEX 2047, neutral lipids (78.6%) were the major component of the total lipids, in which TAG was the predominant component (87.9%) of neutral lipids. EPA was present 37.44% in TAG, 22.49% in MAG and 15.91% in PC.¹³⁸ EPA content increased in polar lipids but decreased in neutral lipids at 10 – 30 g/L NaCl.¹³⁹ When increasing the temperature from 15 to 23 °C, EPA content slightly decreased in TAG but increased in glycolipids. EPA content in phospholipids at 19 °C was higher than that at 15 °C and 23 °C.¹⁴⁰ In *S. mangrovei* FB3, TAG was the predominant component with 97.2% of neutral lipids. Neutral lipids constitute 95.9% of total lipids. PC was the major polar lipids which accounted for 47.78% of phospholipids. DHA was found as the main polyunsaturated fatty acid since it was 29.74% in TAG and 39.61% in PC.⁸⁵ PC in *C. cohnii* was the major component (63.6%) of polar lipids in which 57.2% were DHA.¹⁴¹ However, it was stated that DHA accumulated predominantly in *C. cohnii* cells as TAG, the neutral lipid fraction.¹⁰

N. saprophila was mixotrophically grown with acetate in which PC was the major component (55.7% of lipids) and EPA was concentrated 28.2% in PC whereas PC was only 47.9% of lipids and EPA was 19.0% of PC in photoautotrophic culture.⁷⁴

In the photoautotrophic culture of *P. incisa*, TAG was the dominant lipid with 42.9% of fatty acids in the logarithmic phase and 77% in the stationary phase. ARA was mainly present in TAG with 43% in logarithmic phase and 47% in stationary phase.⁵⁹ Under nitrogen starvation, neutral lipids and ARA content in neutral lipids were 86.8% and 63.7% of total fatty acids compared to 62.1% and 50.8% in the control, respectively.⁶⁰ In *P. lutheri*, TAG was the major component of nonpolar lipids and MGDG was the main component of polar lipids. EPA was present 45% in MGDG and 33% in TAG. DHA was distributed 27% in TAG, 22% in DPG and 21% in betaine lipids.⁸² Light affected distribution of EPA in lipid classes. Under low light intensity (9 W/m²), EPA accumulated in polar lipids was higher than that in non-polar lipids whereas it was conversely when cultured at 19 and 30 W/m². At these conditions of light,

DHA content in non-polar lipids was higher than that in polar lipids. When increasing light intensity from 9 to 30 W/m², EPA and DHA contents in polar lipids decreased while EPA and DHA contents in non-polar lipids at 19 W/m² were higher than those at 9 and 30 W/m².¹³² Galactolipid fraction contained 92% GLA in *S. platensis* 2340.¹⁴² Nitrogen starvation affected distribution of ARA and EPA in *P. cruentum*. ARA content in total fatty acids increased from 19.9% to 30.7% in the neutral lipids and from 46.3% to 61.2% in PC whereas EPA decreased from 43.2% to 16.9% in MGDG, 29.4% to 8.6% in SQ and 17.4% to 2.9% in PC.⁵⁸

Formation of lipid bodies was revealed by using fluorescent staining of endoplasmic reticulum (ER). Lipid bodies surrounded ER in oleaginous fungus *M. ramanniana* IFO 8187.¹⁴³ The same result was observed in *S. limacinum* SR21. The lipid bodies often contact with ER in all stages of the cells.¹⁴⁴

In the photosynthetic microorganisms, the lipid body formation occurred in the inner thylakoid spaces of the chloroplast structure in *Isochrysis*¹⁴⁵ or *M. subterraneus* UTEX 151.⁶⁵

Table 2. Distribution of LC-PUFAs in lipid classes.

Microorganisms	Modes of nutrition	LC-PUFAs	LC-PUFAs (mg/g dry weight)	
			Polar lipids	Non polar lipids
<i>Cunninghamella echinulate</i> ATTHUM 4411 ³² (a)	H	GLA	0.93	7.04
<i>M. alpina</i> SC9 ¹³⁷ (b)	H	ARA	1.22	84.95
<i>N. laevis</i> UTEX 2047 ¹³⁹ (c)	H	EPA	9.81	61.19
<i>N. laevis</i> UTEX 2047 ¹³⁸	H	EPA	5.87	9.11
<i>S. mangrovei</i> FB3 ⁸⁵	H	DHA	9.00	193.17
<i>P. lutheri</i> SMBA 60 ⁸²	P	EPA	8.41*	4.95*
<i>P. lutheri</i> SMBA 60 ⁸²	P	DHA	6.39*	2.26*
<i>P. lutheri</i> SMBA 60 ¹³² (d)	P	EPA	526.94*	170.77*
<i>P. lutheri</i> SMBA 60 ¹³² (d)	P	DHA	69.77*	177.99*

*Unit: mg/g ash free dry weight (AFDW); (a) late exponential phase; (b) 0 % NaCl; (c) 10 g/L NaCl; (d) 9 W/m²

6. LC-PUFA YIELD AND PRODUCTIVITY OF PHOTOSYNTHETIC AND NON-PHOTOSYNTHETIC MICROORGANISMS

Non-photosynthetic microorganisms only grow and produce LC-PUFAs in heterotrophic condition while photosynthetic microorganisms

can grow and produce LC-PUFAs in photoautotrophic, mixotrophic and heterotrophic conditions (Table 4). The LC-PUFA producers for high productivity have been selected to compare the performance of their production (Table 3).

Table 3. Comparison of LC-PUFA yield and productivity of selected microorganisms.

LC-PUFAs	Microorganisms	Modes of nutrition	Biomass (g/L)	LC-PUFA yield (g/L)	LC-PUFA productivity (mg/L/d)
GLA	<i>M. rouxii</i> CBS 416.77 ³⁵	H	24.0	0.532	336.0
	<i>C. echinulata</i> CCRC 31840 ³¹	H	38.1	1.349	269.8
	<i>M. ramanniana</i> CBS 112.08 ²⁷	H	12.0	0.451	112.8
	<i>S. platensis</i> M2 ²⁸	P	-	-	26.4
ARA	<i>M. alpina</i> DSA-12 ¹¹⁴	H	72.5	18.800	1504.0
	<i>M. alpina</i> ME-1 ¹⁴⁶	H	39.8	19.020	3396.4
	<i>P. incisa</i> comb. nov ¹⁴⁷	P	21.0	2.667	70.2
EPA	<i>M. alpina</i> 20-17 ¹⁴⁸	H	24.5	1.350	103.8
	<i>N. laevis</i> UTEX 2047 ⁷⁷	H	-	-	174.6
	<i>P. irregulare</i> ⁸⁴	H	14.22	0.176	-
	<i>P. tricornutum</i> UTEX 640 ⁵⁶	M	15.4	0.436	43.1
	<i>P. tricornutum</i> UTEX 640 ¹⁴⁹	M	-	-	56.0
	<i>P. tricornutum</i> UTEX 640 ⁶⁹	P	1.7	0.083	25.1
	<i>M. subterraneus</i> UTEX 151 ⁶⁶	P	-	-	58.9
DHA	<i>C. cohnii</i> ATCC 30772 ⁸⁷	H	109	11.700	1276.4
	<i>S. limacinum</i> ATCC 1381 ⁴⁸	H	48.1	13.300	3325.0
	<i>Schizochytrium</i> ¹³⁵	H	178	33.286	16560.0
	<i>I. galbana</i> UTEX LB 2307 ¹⁵⁰	P	-	-	4.3

6.1. Heterotrophic production

Until now, numerous data of heterotrophic LC-PUFA production by non-photosynthetic microorganisms have been published.

For GLA production, fungi were found as producers in high GLA productivity. *M. rouxii* CBS 416.77 was cultivated with glucose and Difco yeast extract. Biomass and GLA productivity were 24 g/L and 336 mg/L/d, respectively.³⁵

Higashiyama *et al.* has compared productivity of ARA production by *Mortierella*, in which strain *M. alpina* 1S-4 gave high ARA productivity (1300 mg/L/d).³⁶ However, Hwang *et al.* cultivated *M. alpina* DSA-12 in fed-batch

by using NH₄OH as a nitrogen source and pH control which obtained 1504 mg/L/d ARA, higher than former productivity in the culture of Higashiyama *et al.*¹¹⁴ *M. alpina* ME-1 was a UV-mutant of ATCC 16266 gave 19020 mg/L ARA at 5.6 days which was highest found in the reports.¹⁴⁶

EPA production has been reviewed by Bajpai and Bajpai.¹⁵ High EPA yield and productivity were 1350 mg/L and 103.8 mg/L/d in the culture of *M. alpina* 20-17.¹⁴⁸ *N. laevis* UTEX 2047 was grown in perfusion culture with cell bleeding. EPA productivity obtained in this cultivation (174.6 mg/L/d) was highest EPA productivity found.⁷⁷

Stramenopiles were utilized as DHA producing microorganisms. DHA yield and productivity have been compared among various strains. *S. limacinum* SR21 gave the highest DHA yield and productivity with 13300 mg/L and 138 mg/L/h DHA.^{151,152} The other strain of *Schizochytrium* which has been studied by Bailey *et al.* produced a very high concentration of DHA 23.45 g/L in 42 h.¹³⁵ *C. cohnii* was also a DHA producing microalga. The parameters of culture and DHA production were collected in the review of Mendes *et al.*¹⁰ The fed-batch cultivation on ethanol produced 11700 mg/L DHA in 220 h was the highest productivity in this microalga.⁸⁷

6.2. Mixotrophic production

Mixotrophic production was found in photoautotrophic microorganisms which have growth capacity with organic compounds under light. Up to now, most of LC-PUFA producing microorganisms in mixotrophic condition were EPA producing microalgae and GLA producing cyanobacteria (Table 4). *P. tricornutum* UTEX-640 was cultivated with carbon sources to evaluate growth and EPA production. Glycerol was found as the most suitable source of carbon.^{56,69} Fed-batch culture with 0.1 M concentration of glycerol and the successive additions of ammonium chloride gave 16.2 g/L biomass concentration, 61.5 mg/L/h biomass productivity and 33.5 mg/L/d EPA productivity. This EPA productivity was 10-fold greater than the maximum productivity obtained in the

photoautotrophic control culture.⁵⁵ Additionally, fed-batch with glycerol and urea gave 43.13 mg/L/d EPA productivity which was 13-fold higher than the maximum EPA productivity obtained in photoautotrophic culture of the control.⁵⁶ Other result of *P. tricornutum* UTEX-640 indicated that EPA productivity (56 mg/L/d) in mixotrophic culture (with glycerol) was approximately 3-fold higher than that in photoautotrophic culture.¹⁴⁹ *N. saprophila* was mixotrophically cultivated with acetate. EPA content obtained in this condition was 19.2 mg/g biomass that was higher than those obtained in photoautotrophic and heterotrophic conditions.⁷³ EPA content in biomass was 34.6 mg/g when *N. saprophila* was cultured with 2 mM acetate and 2% CO₂.⁷⁴ Performance of EPA production in three nutritional modes was compared in the culture of *N. laevis* UTEX 2047. Growth and EPA production were highest in mixotrophic culture. EPA yield and productivity were 52.32 mg/L and 10.46 mg/L/d, respectively.¹³ *Nannochloropsis* sp. also showed that they can grow and produce EPA in 3 nutritional modes. Glucose and ethanol were utilized as carbon sources for EPA production which gave 23.4 mg/L and 23.0 mg/L EPA, respectively, in mixotrophic cultivation after 8 days.⁷² An increase of EPA yield up to 56 mg/L was obtained in 10 days of fed-batch culture with an addition of glucose and nitrate.¹⁵² *S. platensis* KCTC AG20590 was mixotrophically cultivated with the long or short chain carbon sources. Results indicated that GLA content increased when compared with the control.⁸³

Table 4. Comparison of LC-PUFA productivity between the nutritional modes.

Strains	LC-PUFAs	LC-PUFA productivity (mg/L/d)		
		PA	M	H
<i>Nannochloropsis</i> sp. ⁷²	EPA	3.13	3.34	1.44
<i>N. laevis</i> UTEX 2047 ¹³	EPA	3.39	10.46	6.37
<i>P. tricornutum</i> UTEX-640 ⁵⁶	EPA	3.35	43.13	-
<i>P. tricornutum</i> UTEX-640 ¹⁴⁹	EPA	18.0	56.0	-
<i>N. saprophila</i> ⁷⁴	EPA	4.93	14.8	-
<i>S. platensis</i> KCTC AG20590 ⁸³	GLA	0.43	1.70	-

6.3. Photoautotrophic production

Photoautotrophic production was only found in photosynthetic microorganisms. Table 5 presented performance of essential LC-PUFAs from photosynthetic microorganisms.

GLA productivity was 26.4 mg/L/d obtained from *S. platensis* M2 in outdoor culture.⁶⁶

*Parietochloris*¹⁴⁷ and *Porphyridium*¹⁵³ were ARA-producing microalgae. EPA productivity was 70.2 mg/L/d and 6.5 mg/L/d, respectively.

For EPA production, *Phaeodactylum* was known as photoautotrophic EPA producer. Meiser *et al.* cultivated *P. tricornutum* UTEX

640 under continuous light in batch culture.¹⁵⁴ Maximal EPA productivity 118 mg/L/d were obtained. *Nannochloropsis* sp. was cultivated in flat plate reactor under 1000 μ mol photon/m²/s gave 127.9 mg/L/d EPA.⁷¹

Until now, *Rhodomonas*, *Pavlova* and *Isochrysis* were found as photosynthetic microalgae produced DHA. However, productivity of DHA production by these microalgae was less than that by non-photosynthetic microorganisms (Table 3). The highest DHA productivity was 4.3 mg/L/d, obtained when cultured *I. galbana* in optical fiber photobioreactor.¹⁵⁰

Table 5. LC-PUFA productivity of photoautotrophic production.

LC-PUFAs	Strains	LC-PUFA productivity (mg/L/d)
GLA	<i>S. platensis</i> M2 ⁶⁶	26.4
	<i>S. platensis</i> KCTC AG20590 ⁸³	0.4
ARA	<i>P. incisa</i> comb. nov ¹⁴⁷	70.2
	<i>P. cruentum</i> IAM R-3 ¹⁵³	6.5
EPA	<i>S. putrefaciens</i> MAC1 ²⁴	58.3
	<i>P. cruentum</i> IAM R-3 ¹⁵³	3.6
	<i>M. subterraneus</i> UTEX 151 ¹⁵⁴	56.0
	<i>M. subterraneus</i> UTEX 151 ⁶⁴	25.7
	<i>M. subterraneus</i> UTEX 151 ⁶⁶	58.9
	<i>P. tricornutum</i> UTEX 640 ⁶⁸	19.0
	<i>P. tricornutum</i> UTEX 640 ⁶⁹	25.1
	<i>P. tricornutum</i> UTEX 640 ¹⁵⁴	50.0
	<i>P. tricornutum</i> UTEX 640 ¹⁵⁵	13.0
	<i>P. tricornutum</i> UTEX 640 ¹⁵⁶	118.0
	<i>P. tricornutum</i> UTEX 640 ¹⁵⁷	47.8
	<i>P. tricornutum</i> TFX-1 ¹⁵⁸	6.0
	<i>Nannochloropsis</i> sp. ⁷⁰	32.0
	<i>Nannochloropsis</i> sp. ⁷¹	127.9
	<i>Nannochloropsis</i> sp. PP983 ¹¹¹	1.2
	<i>P. lutheri</i> SMBA 60 ¹⁶⁰	0.5
	<i>P. lutheri</i> SMBA 60 ¹³²	1.3
	<i>I. galbana</i> Parke ¹⁶⁰	4.8
	<i>I. galbana</i> ¹⁶¹	15.3
	<i>I. galbana</i> ¹⁶²	7.2
DHA	<i>P. lutheri</i> SMBA 60 ¹⁵⁹	0.2
	<i>P. lutheri</i> SMBA 60 ¹³²	0.7
	<i>I. galbana</i> UTEX LB 2307 ¹⁵¹	4.3
	<i>I. galbana</i> CCMP 1324 ¹⁶³	0.6
	<i>I. galbana</i> CCAP 927/1 ¹⁶⁴	0.2
	<i>I. galbana</i> ¹⁶²	3.1

7. YIELD CONVERSION OF LC-PUFA PRODUCTION

Conversion yield is calculated on the ratio of production to substrate. It permits to evaluate

Table 6. Biomass and LC-PUFAs conversion yield of microorganisms.

Microorganisms	Substrate (S)	Y_{XS} (g. g ⁻¹)	LC-PUFAs	$Y_{LC-PUFA/S}$ (mg. g ⁻¹)	Modes of nutrition
<i>M. circinelloides</i> CBS 203.28 ³⁴	Acetic acid	0.30	GLA	10.00	H
<i>C. cohnii</i> ATCC 30772 ¹⁰⁸	Acetic acid	0.12	DHA	30.00	H
<i>M. isabellina</i> ATHUM 2935 ³⁵	Glucose	0.50	GLA	7.70	H
<i>M. alpina</i> DSA-12 ¹¹⁴	Glucose	0.44	ARA	95.40	H
<i>Schizochytrium</i> G13/2S ¹⁶⁵	Glucose	0.39	DHA	64.02	H
<i>Aurantiochytrium limacinum</i> mh0186 ⁵²	Glucose	0.38	DHA	71.67	H
Thraustochytrid G13 ¹⁶⁶	Glucose	0.46	DHA	56.25	H
<i>C. cohnii</i> ATCC 30556 ¹²⁸	Glucose	0.50	DHA	38.59	H
<i>C. cohnii</i> ATCC 30772 ⁸⁶	Glucose	0.37	DHA	21.27	H
<i>Nannochloropsis</i> sp. ⁷²	Glucose	0.20	EPA	8.75	M
<i>N. laevis</i> UTEX 2047 ¹³	Glucose	0.42	EPA	10.46	M

X: biomass; Y: conversion yield

8. IMPROVEMENT FOR LC-PUFA PRODUCTION

Microalgae cultivation in large volume increases the productivity of biomass and LC-PUFAs. Nowadays, a lot of photobioreactors were invented for microalgae culture.

Cultivation of non-photosynthetic microorganisms was carried out in the closed and sterile systems with the sources of organic carbon. Because of heterotrophic culture, light was not necessary in this system. Source of carbon is usually one of the factors influencing production. Thus, fed-batch or continuous culture were often used to improve LC-PUFA production.

Conversely, light was necessary in cultivation of photosynthetic microorganisms.¹⁶⁷ Thus, bioreactors can be designed to obtain light effectively. By using a new type of enclosed photobioreactor in which light was efficiently distributed by light diffusing optical fibers, DHA

productivity economically. Table 6 showed conversion yield of biomass and LC-PUFAs in some LC-PUFA producers.

from *Isochrysis* was obtained 4.3 mg/L/d (Table 3), twofold greater than that obtained using flat glass bottles.¹⁵⁰ *Nannochloropsis* sp. was cultured in a flat plate reactor with a narrow (1 – 2 cm) light path and rigorous stirring exposed to high photon flux densities (1000-3000 μ mol photons/m²/s). Biomass and EPA yield were obtained 40.6 g/L and 2302 mg/L, respectively.⁷¹

Culture in two stages of temperature is a strategy for improvement of LC-PUFA production: the first stage for biomass production and the second for LC-PUFA production. In the second stage, temperature was usually decreased to produce more LC-PUFAs.

In *P. irregulare* ATCC 10951, cells were initially grown at 25 °C for 1, 2 or 3 days and then shifted to 12 °C for 6, 8, 9 days. The best combination was 2 days at 25 °C, followed by 6 days at 12 °C, which gave 93.1 μ g/ml EPA.³⁹ *M. alpina* ATCC 32222 was cultured for 8 days at 25 °C gave a high biomass (52.4 g/L)

and ARA yield (9.1 g/L). Then, the culture was incubated at 15 °C. The maximal ARA content was obtained (11.1 g/L) in 11 days of fermentation.¹⁶⁸ An increase in cellular DHA content by 19.9% and productivity by 6.5% was observed when the temperature in the culture of *C. cohnii* ATCC 30556 was shifted from 25 °C for 2 days to 15 °C for 1 day compared with that maintained at 25 °C for 3 days.¹²⁶ A shift of temperature from 30 °C for 32 h to 20 °C for 12 h in the culture of *Schizochytrium* sp. HX-308 resulted in an increase of DHA content which is present 6.05% in dry cell weight and 51.98% in total fatty acids.¹⁶⁹ In *C. cohnii* CCMP 316, n-dodecane was added in the culture as an oxygen vector. The DHA content in total fatty acids, the DHA content in biomass and DHA yield increased by 16, 39 and 22%, respectively, at 0.5% n-dodecane.⁵³

The increase of DHA content (15.7 to 17.8 mg/g biomass) was also found when *I. galbana* LB 2307 was shifted from 24 °C to 17 °C for 24 h.¹⁵⁰ In the culture of *P. tricornutum* 2038, cells were cultivated at 25 °C and then shifted to 20, 15, 10 °C. An increase of EPA content per dry mass was observed after 12 h, 24 h and 48 h at 10 °C, 15 °C and 20 °C, respectively. The highest EPA yield was 6.6 mg/L when temperature was shifted from 25 °C to 10 °C for 12 h, which raised by 120% compared with the control.¹²³ After decantation, biomass of *S. costatum* was obtained and incubated at 15 °C for 15 h, resulted in an increase in EPA content from 11 mg/g to 19 mg/g of dry weight.⁷⁹

9. CONCLUSION

The limitation of essential LC-PUFA sources originating from animals and plants has promoted the research on other sources. Microorganisms were found as potential sources for LC-PUFA production because they could grow fast on culture media and contain high LC-PUFA content in their cells. Besides heterotrophically LC-PUFA producing microorganisms, many microalgae have been discovered as LC-PUFA producers. Among

these microalgae, some strains could produce LC-PUFAs in 2 or 3 modes of nutrition. Among nutritional modes, heterotrophy was found as a mode of high productivity production. However, mixotrophy has also potential for improvement of LC-PUFA productivity in photosynthetic microorganisms. Further researches need to focus on new microalgal strains to diversify LC-PUFA sources.

REFERENCES

1. O. P. Ward, A. Singh. Omega-3/6 fatty acids: alternative sources of productions, *Process Biochemistry*, **2005**, *40*, 3267-3652.
2. C. H. Ruxton, S. C. Reed, M. J. Simpson, K. J. Millington. The health benefits of omega-3 polyunsaturated fatty acids: a review of the evidence, *Journal of Human Nutrition and Dietetics*, **2004**, *17*, 449-459.
3. A. Molfino, M. I. Amabile, M. Monti, S. Arcieri, F. F. Rossi, M. Muscaritoli. The role of docosahexaenoic acid (DHA) in the control of obesity and metabolic derangements in breast cancer, *International Journal of Molecular Science*, **2016**, *17*, 505.
4. K. Christina, F. Maria, S. Anastasis, P. Ioannis. DHA/EPA (Omega-3) and LA/GLA (Omega-6) as bioactive molecules in neurodegenerative diseases, *International Journal of Molecular Sciences*, **2023**, *24*, 10717.
5. W. S. Harris. Fish oil supplementation: evidence for health benefits, *Cleveland Clinic Journal of Medicine*, **2004**, *71*, 209-221.
6. U. Varanasi, J. E. Stein. Disposition of xenobiotic chemicals and metabolites in marine organisms, *Environmental Health Perspectives*, **1991**, *90*, 93-100.
7. J. S. Christiansen, R. A. Dalmo, K. Ingebrigtsen. Xenobiotic excretion in fish with agglomerular kidneys, *Marine Ecology Progress Series*, **1996**, *136*, 303-304.
8. N. T. Eriksen. The technology of microalgal culturing. *Biotechnology Letters*, **2008**, *30*, 1525-1536.

9. C. U. Ugwu, H. Aoyagi, H. Uchiyama. Photobioreactors for mass cultivation of algae, *Bioresource Technology*, **2008**, 99, 4021-4028.
10. A. Mendes, A. Reis, R. Vasconcelos, P. Guerra, T. L. D. Silva. Cryptocodinium cohnii with emphasis on DHA production: a review, *Journal of Applied Phycology*, **2009**, 21, 199-214.
11. W. R. Barclay, K. M. Meager, J. R. Abril. Heterotrophic production of long-chain omega-3 fatty acids utilizing algae and algae-like microorganisms, *Journal of Applied Phycology*, **1994**, 6, 123-129.
12. I. K. Goldberg, U. Iskandarov, Z. Cohen. LC-PUFA from photosynthetic microalgae: occurrence, biosynthesis, and prospects in biotechnology, *Applied Microbiology and Biotechnology*, **2011**, 91, 905-915.
13. Z. Y. Wen, F. Chen. Production potential of eicosapentaenoic acid by the diatom *nitzschia laevis*, *Biotechnology Letters*, **2000**, 22, 727-733.
14. U. Bandyopadhyay, D. Das, R. K. Banerjee. Reactive oxygen species: oxidative damage and pathogenesis, *Current Science*, **1999**, 77, 658-666.
15. P. Bajpai, P. K. Bajpai. Eicosapentaenoic acid (EPA) production from microorganisms: a review, *Journal of Biotechnology*, **1993**, 30, 161-183.
16. Z. Y. Wen, F. Chen. Heterotrophic production of eicosapentaenoic acid by microalgae, *Biotechnology Advances*, **2003**, 21, 273-294.
17. D. J. Kyle, S. E. Reeb, V. J. Sicotte. Production of docosahexaenoic acid by dinoflagellates, United States Patent, US 5407957, 1995.
18. A. L. Reysenbach, S. L. Cady. Microbiology of ancient and modern hydrothermal systems, *Trends in Microbiology*, **2001**, 9, 79-86.
19. L. Xu, P. J. Weathers, X. R. Xiong, C. Z. Liu. Microalgal bioreactors: challenges and opportunities, *Engineering in Life Sciences*, **2009**, 9, 178-189.
20. B. Wang, C. Q. Lan, M. Horsman. Closed photobioreactors for production of microalgal biomasses, *Biotechnology Advances*, **2012**, 30, 904-912.
21. D. S. Nichols, J. L. Brown, P. D. Nichols, T. A. McMeekin. Production of eicosapentaenoic acid and arachidonic acids by an antarctic bacterium: response to growth temperature, *FEMS Microbiology Letters*, **1997**, 152, 349-354.
22. K. Watanabe, C. Ishikawa, I. Ohtsuka, M. Kamata, M. Tomita, K. Yazawa, H. Muramatsu. Lipid and fatty acid compositions of a novel docosahexaenoic acid-producing marine bacteria, *Lipids*, **1997**, 32, 975-978.
23. Y. S. Jeong, S. K. Song, S. J. Lee, B. K. Hur. The growth and EPA synthesis of *shewanella oneidensis* MR-1 and expectation of EPA biosynthetic pathway, *Biotechnology and Bioprocess Engineering*, **2006**, 11, 127-133.
24. N. Houshang, G. Mansel. Production of EPA by *shewanella putrefaciens* MAC1 in selected culture media, *Journal of Food Science and Technology*, **2008**, 5, 75-81.
25. E. D. Frank, E. M. Joseph, J. J. Brittany, S. Sydney, P. L. James, J. D. Kellen, P. R. Michael, B. M. James. The microbiota of freshwater fish and freshwater niches contain omega-3 fatty acid-producing *shewanella* species, *Applied and Environmental Microbiology*, **2016**, 82, 218-231.
26. N. Morita, M. Tanaka, H. Okuyama. Biosynthesis of fatty acids in the docosahexaenoic acid-producing bacterium *moritella marina* strain MP-1, *Biochemical Society Transactions*, **2000**, 28, 943-945.
27. A. M. Lindberg, G. Molin. Effect of temperature and glucose supply on the production of polyunsaturated fatty acids by the fungus *mortierella alpina* CBS 343.66 in fermentor cultures, *Applied Microbiology and Biotechnology*, **1993**, 39, 450-455.
28. O. Hiruta, Y. Kamisaka, T. Yokochi, T. Futamura, H. Takebe, A. Satoh, T. Nakahara, O. Suzuk. Gamma-linoleic acid production by a low temperature-resistant mutant of *mortierella ramanniana*, *Journal of Fermentation and Bioengineering*, **1996**, 82, 119-123.
29. M. Xian, J. Nie, Q. Meng, J. Liu, C. Zhou, Y. Kang, K. Zhen. Production of γ -linolenic acid by disrupted mycelia of *mortierella isabellina*,

Letters in Applied Microbiology, **2003**, *36*, 182-185.

30. H. C. Chen, C. C. Chang. Production of γ -linolenic acid by the fungus *cunninghamella echinulata* CCRC 31840, *Biotechnology Progress*, **1996**, *12*, 338-341.

31. H. C. Chen, T. M. Liu. Inoculum effects on the production of γ -linolenic acid by the shake culture of *cunninghamella echinulata* CCRC 31840, *Enzyme and Microbial Technology*, **1997**, *21*, 137-142.

32. H. Gema, A. Kavadia, D. Dimou, V. Tsagou, M. Komaitis, G. Aggelis. Production of γ -linolenic acid by *cunninghamella echinulata* cultivated on glucose and orange peel, *Applied Microbiology and Biotechnology*, **2002**, *58*, 303-307.

33. K. C. Naveena, Ramalingappa. Production of polyunsaturated fatty acids (PUFAs) from microbes and their secondary metabolites, *International Journal of Current Microbiology Applied Sciences*, **2018**, *7*, 2680-2689.

34. A. M. Lindberg, L. Hansson. Production of γ -linolenic acid by the fungus *mucor rouxii* on cheap nitrogen and carbon sources, *Applied Microbiology and Biotechnology*, **1991**, *36*, 26-28.

35. E. V. Emelyanova. Lipid and γ -linolenic acid production by *mucor inaquisporus*, *Process Biochemistry*, **1997**, *32*, 173-177.

36. P. Bajpai, P. Bajpai, O. P. Ward. Arachidonic acid production by fungi, *Applied and Environmental Microbiology*, **1991**, *57*, 1255-1258.

37. K. Higashiyama, S. Fujikawa, E. Y. Park, S. Shimizu. Production of arachidonic acid by *mortierella* fungi, *Biotechnology and Bioprocess Engineering*, **2002**, *7*, 252-262.

38. S. R. Gandhi, J. D. Weete. Production of the polyunsaturated fatty acids arachidonic acid and eicosapentaenoic acid by the fungus *pythiumultimum*, *Journal of General Microbiology*, **1991**, *137*, 1825-1830.

39. E. E. Stinson, R. Kwoczak, M. J. Kurantz. Effect of cultural conditions on production of eicosapentaenoic acid by *pythium irregulare*, *Journal of Industrial Microbiology*, **1991**, *8*, 171-178.

40. P. Bajpai, P. K. Bajpai, O. P. Ward. Eicosapentaenoic acid (EPA) production by *mortierella alpina* ATCC 32222, *Applied Biochemistry and Biotechnology*, **1991**, *31*, 267-272.

41. P. K. Bajpai, P. Bajpai, O. P. Ward. Optimisation of culture conditions for production of eicosapentaenoic acid by *mortierella elongata* NRRL 5513, *Journal of Industrial Microbiology*, **1992**, *9*, 11-18.

42. L. Bongiorni, R. Jain, S. Raghukumar, R. Aggarwal. *Thraustochytrium gaertnerium* sp. nov.: a new thraustochytrid stramenopilan protist from mangroves of Goa, India, *Protistología*, **2005**, *156*, 303-315.

43. A. Singh, S. Wilson, O. P. Ward. Docosahexaenoic acid (DHA) production by *thraustochytrium* sp. ATCC 20892, *World Journal of Microbiology and Biotechnology*, **1996**, *12*, 76-81.

44. P. K. Bajpai, P. Bajpai, O. P. Ward. Optimization of production of docosahexaenoic acid (DHA) by *thraustochytrium aureum* ATCC 34304, *Journal of the American Oil Chemists' Society*, **1991**, *68*, 509-514.

45. Z. Y. Li, O. P. Ward. Production of docosahexaenoic acid by *thraustochytrium roseum*, *Journal of Industrial Microbiology*, **1994**, *13*, 238-241.

46. A. Singh, O. P. Ward. Production of high yields of docosahexaenoic acid by *thraustochytrium roseum* ATCC 28210, *Journal of Industrial Microbiology*, **1996**, *16*, 370-373.

47. A. M. Burja, H. Radianingtyas, A. Windust, C. J. Barrow. Isolation and characterization of polyunsaturated fatty acid producing *thraustochytrium* species: screening of strains and optimization of omega-3 production, *Applied Microbiology and Biotechnology*, **2006**, *72*, 1161-1169.

48. T. Yaguchi, S. Tanaka, T. Yokochi, T. Nakahara, T. Higashihara. Production of high yields of docosahexaenoic acid by *schizochytrium* sp. strain SR21, *Journal of the American Oil Chemists' Society*, **1997**, *74*, 1431-1434.

49. T. Yokochi, D. Honda, T. Higashihara, T. Nakahara. Optimization of docosahexaenoic

acid production by *schizochytrium limacinum* SR21, *Applied Microbiology and Biotechnology*, **1998**, *49*, 72-76.

50. Z. Chi, Y. Liu, C. Frear, S. Chen. Study of a two-stage growth of DHA-producing marine algae *schizochytrium limacinum* SR21 with shifting dissolved oxygen level, *Journal of Applied Phycology*, **2009**, *81*, 1141-1148.

51. M. K. M. Wong, C. M. K. Tsui, D. W. T. Au, L. L. P. Vrijmoed. Docosahexaenoic acid production and ultrastructure of the thraustochytrid *aurantiochytrium mangrovei* MP2 under high glucose concentrations, *Mycoscience*, **2008**, *49*, 266-270.

52. N. Nagano, Y. Taoka, D. Honda, M. Hayashi. Optimization of culture conditions for growth and docosahexaenoic acid production by a marine thraustochytrid, *aurantiochytrium limacinum* mh0186, *Journal of Oleo Science*, **2009**, *58*, 623-628.

53. T. L. D. Silva, A. Reis. The use of multiparameter flow cytometry to study the impact of n-dodecane additions to marine dinoflagellate microalga *cryptocodinium cohnii* batch fermentations and DHA production, *Journal of Industrial Microbiology and Biotechnology*, **2008**, *35*, 875-887.

54. C. K. Tan, M. Johns. Screening of diatoms for heterotrophic eicosapentaenoic acid production, *Journal of Applied Phycology*, **1996**, *8*, 59-64.

55. M. C. C. García, J. M. F. Sevilla, F. G. A. Fernández, E. M. Grima, F. G. Camacho. Mixotrophic growth of *phaeodactylum tricornutum* on glycerol: growth rate and fatty acid profile, *Journal of Applied Phycology*, **2000**, *12*, 239-248.

56. M. C. C. García, A. S. Mirón, J. M. F. Sevilla, E. M. Grima, F. G. Camacho. Mixotrophic growth of the microalga *phaeodactylum tricornutum*, influence of different nitrogen and organic carbon sources on productivity and biomass composition, *Process Biochemistry*, **2005**, *40*, 297-305.

57. M. Mühling, A. Belay, A. B. Whitton. Variation in fatty acid composition of *arthrospira* (*spirulina*) strains, *Journal of Applied Phycology*, **2005**, *17*, 137-146.

58. Z. Cohen. The production potential of eicosapentaenoic and arachidonic acids by the red alga *porphyridium cruentum*, *Journal of the American Oil Chemists' Society*, **1990**, *67*, 916-920.

59. C. Bigogno, I. K. Goldberg, S. Boussiba, A. Vonshak, Z. Cohen. Lipid and fatty acid composition of the green oleaginous alga *parietochloris incisa*, the richest plant source of arachidonic acid, *Phytochemistry*, **2002**, *60*, 497-503.

60. I. K. Goldberg, C. Bigogno, P. Shrestha, Z. Cohen. Nitrogen starvation induces the accumulation of arachidonic acid in the freshwater green alga *parietochloris incisa* (trebouxiophyceae), *Journal of Phycology*, **2002**, *38*, 991-994.

61. A. E. Solovchenko, I. K. Goldberg, S. D. Cohen, Z. Cohen, M. N. Merzlyak. Effects of light intensity and nitrogen starvation on growth, total fatty acids and arachidonic acid in the green microalga *parietochloris incisa*, *Journal of Applied Phycology*, **2008**, *20*, 245-251.

62. V. L. Angélica, A. Felipe, N. Karla. Microalgae, a potential natural functional food source – a review, *Polish Journal of Food and Nutrition Sciences*, **2017**, *67*, 251-263.

63. G. López, C. Yate, F. A. Ramos, M. P. Cala, S. Restrepo, S. Baena. Production of polyunsaturated fatty acids and lipids from autotrophic, mixotrophic and heterotrophic cultivation of *galdieria* sp. strain USBA- GBX-832, *Scientific Reports*, **2019**, *9*, 10791.

64. Z. Cohen. Production potential of eicosapentaenoic acid by *monodus subterraneus*, *Journal of the American Oil Chemists' Society*, **1994**, *71*, 941-945.

65. H. Qiang, H. Zhengyu, Z. Cohen, A. Richmond. Enhancement of eicosapentaenoic acid (EPA) and γ -linolenic acid (GLA) production by manipulating algal density of outdoor cultures of *monodus subterraneus* (eustigmatophyta) and *spirulina platensis* (Cyanobacteria), *European Journal of Phycology*, **1997**, *32*, 81-86.

66. C. P. Liu, L. P. Lin. Morphology and eicosapentaenoic acid production by *monodus subterraneus* UTEX 151, *Micron*, **2005**, *36*, 545-550.

67. Z. Cohen, S. Didi, Y. M. Heimer. Overproduction of γ -linolenic and eicosapentaenoic acids by algae, *Plant Physiology*, **1992**, *98*, 569-572.

68. W. Yongmanitchai, O. P. Ward. Growth of and omega-3 fatty acid production by *phaeodactylum tricornutum* under different culture conditions, *Applied Microbiology and Biotechnology*, **1991**, *57*, 419-425.

69. W. Yongmanitchai, O. P. Ward. Growth and eicosapentaenoic acid production by *phaeodactylum tricornutum* in batch and continuous culture systems, *Journal of the American Oil Chemists' Society*, **1992**, *69*, 584-590.

70. G. C. Zittelli, F. Lavista, A. Bastianini, L. Rodolfi, M. Vincenzini, M. R. Tredici. Production of eicosapentaenoic acid by *nannochloropsis* sp. cultures in outdoor tubular photobioreactors, *Journal of Biotechnology*, **1999**, *70*, 299-312.

71. N. Zou, C. Zhang, Z. Cohen, A. Richmond. Production of cell mass and eicosapentaenoic acid (EPA) in ultrahigh cell density cultures of *nannochloropsis* sp. (Eustigmatophyceae), *European Journal of Phycology*, **2000**, *35*, 127-133.

72. F. Xu, W. Cong, Z. L. Cai, Ouyang. Effects of organic carbon sources on cell growth and eicosapentaenoic acid content of *nannochloropsis* sp. *Journal of Applied Phycology*, **2004**, *16*, 499-503.

73. M. Kitano, R. Matsukawa, I. Karube. Changes in eicosapentaenoic acid content of *navicula saprophila*, *rhodomonas salina* and *nitzschia* sp. under mixotrophic conditions, *Journal of Applied Phycology*, **1997**, *9*, 559-563.

74. M. Kitano, R. Matsukawa, I. Karube. Enhanced eicosapentaenoic acid production by *navicula saprophila*, *Journal of Applied Phycology*, **1998**, *10*, 101-105.

75. Z. Y. Wen, F. Chen. Heterotrophic production of eicosapentaenoic acid by the diatom *nitzschia laevis*: effects of silicate and glucose, *Journal of Industrial Microbiology and Biotechnology*, **2000**, *25*, 218-224.

76. Z. Y. Wen, F. Chen. Optimization of nitrogen sources for heterotrophic production of eicosapentaenoic acid by the diatom *nitzschia laevis*, *Enzyme and Microbial Technology*, **2001**, *29*, 341-347.

77. Z. Y. Wen, F. Chen. A perfusion-cell bleeding culture strategy for enhancing the productivity of eicosapentaenoic acid by *nitzschia laevis*, *Applied Microbiology and Biotechnology*, **2001**, *57*, 316-322.

78. Z. Y. Wen, Y. Jiang, F. Chen. High cell density culture of the diatom *nitzschia laevis* for eicosapentaenoic acid production: fed-batch development, *Process Biochemistry*, **2002**, *37*, 1447-1453.

79. A. Blachemain, D. Grizeau. Increased production of eicosapentaenoic acid by *skeletonema costatum* cells after decantation at low temperature, *Biotechnology Techniques*, **1999**, *13*, 497-501.

80. L. Christos, W. Eric, M. Tanja, C. Bert, R. Johan, V. R. Sandra, B. Leen, V. H. Jasper, R. T. Klaas. Effect of pH on *rhodomonas salina* growth, biochemical composition, and taste, produced in semi-large scale under sunlight conditions, *Journal of Applied Phycology*, **2022**, *34*, 1215-1226.

81. R. Vazhappilly, F. Chen. Eicosapentaenoic acid and docosahexaenoic acid production potential of microalgae and their heterotrophic growth, *Journal of the American Oil Chemists' Society*, **1998**, *75*, 393-397.

82. L. A. Meireles, A. C. Guedes, F. X. Malcata. Lipid class composition of the microalga *pavlova lutheri*: eicosapentaenoic and docosahexaenoic acids, *Journal of Agricultural and Food Chemistry*, **2003**, *51*, 2237-2241.

83. E. K. Kim, G. G. Choi, H. S. Kim, C. Y. Ahn, H. M. Oh. Increasing γ -linolenic acid content in *spirulina platensis* using fatty acid supplement and light-dark illumination, *Journal of Applied Phycology*, **2012**, *24*, 743-750.

84. L. R. Giovanni, L. L. Antonio, M. G. Beatriz, V. Vito, R. Raffaele, S. Raffaele, M. Paolo. New Biotechnological production of EPA by *pythium irregulare* using alternative sustainable media obtained from food industry by-products and waste, *Sustainability*, **2023**, *15*, 1147-1162.

85. K. W. Fan, Y. Jiang, Y. W. Faan, F. Chen. Lipid characterization of mangrove thraustochytrid - schizochytrium mangrovei, *Journal of Agricultural and Food Chemistry*, **2007**, 55, 2906-2910.
86. M. E. D. Swaaf, T. C. D. Rijk, G. Eggink, L. Sijtsma. Optimisation of docosahexaenoic acid production in batch cultivations by cryptocodonium cohnii, *Journal of Biotechnology*, **1999**, 70, 185-192.
87. M. E. D. Swaaf, J. T. Pronk, L. Sijtsma. Fed-batch cultivation of the docosahexaenoic-acid-producing marine alga cryptocodonium cohnii on ethanol, *Applied Microbiology and Biotechnology*, **2003**, 61, 40-43.
88. A. Chalimaa, A. Hatzidakia, A. Karnaouri, E. Topakas. Integration of a dark fermentation effluent in a microalgal-based biorefinery for the production of high-added value omega-3 fatty acids, *Applied Energy*, **2019**, 152, 102-109.
89. X. Qiu. Biosynthesis of docosahexaenoic acid (DHA, 22:6-4, 7, 10, 13, 16, 19): two distinct pathways, *Prostaglandins Leukotrienes and Essential Fatty Acids*, **2003**, 68, 181-186.
90. J. G. Metz, P. Roessler, D. Facciotti, C. Levering, F. Dittrich, M. Lassner, R. Valentine, K. Lardizabal, F. Domergue, A. Yamada. Production of polyunsaturated fatty acids by polyketide synthase in both prokaryotes and eukaryotes, *Science*, **2001**, 293, 290-293.
91. J. C. Lippmeier, K. S. Crawford, C. B. Owen, A. Rivas, J. G. Metz, K. E. Apt. Characterization of both polyunsaturated fatty acid biosynthetic pathways in schizochytrium sp, *Lipids*, **2009**, 44, 621-630.
92. M. E. D. Swaaf, T. C. D. Rijk, P. V. D. Meer, G. Eggink, L. Sijtsma. Analysis of docosahexaenoic acid biosynthesis in cryptocodonium cohnii by ¹³C labelling and desaturase inhibitor experiments, *Journal of Biotechnology*, **2003**, 103, 21-29.
93. C. Ratledge. Fatty acid biosynthesis in microorganisms being used for single cell oil production, *Biochimie*, **2004**, 86, 807-815.
94. J. G. Jaworski. Biosynthesis of monoenoic and polyenoic fatty acids. *The biochemistry of plants*, Academic Press, Orlando, 1987.
95. B. Behrouzian, P. H. Buist. Mechanism of fatty acid desaturation: a bioorganic perspective, *Prostaglandins, Leukotrienes and Essential Fatty Acids*, **2003**, 68, 107-112.
96. D. U. Antonio. Biosynthesis of polyunsaturated fatty acids in lower eukaryotes, *Life*, **2006**, 58, 563-571.
97. D. Shiran, I. Khozin, Y. M. Heimer, Z. Cohen. Biosynthesis of eicosapentaenoic acid in the microalga porphyridium cruentum: the use of externally applied fatty acids, *Lipids*, **1996**, 31, 1277-1282.
98. T. Arao, M. Yamada. Biosynthesis of polyunsaturated fatty acids in the marine diatom, *Phaeodactylum Tricornutum*, *Phytochemistry*, **1994**, 35, 1177-1181.
99. C. Bigogno, I. K. Goldberg, D. Adlerstein, Z. Cohen. Biosynthesis of arachidonic acid in the oleaginous microalga parietochloris incisa (chlorophyceae): radiolabeling studies, *Lipids*, **2002**, 37, 209-216.
100. B. Qi, F. Beaudoin, T. Fraser, A. K. Stobart, J. A. Napier, C. M. Lazarus. Identification of a cDNA encoding a novel C18- Δ^9 polyunsaturated fatty acid-specific elongating activity from the docosahexaenoic acid (DHA)-producing microalga isochrysis galbana, *FEBS Letters*, **2002**, 510, 159-165.
101. S. L. Pereira, A. E. Leonard, Y. S. Huang, L. T. Chuang, P. Mukerji. Identification of two novel microalgal enzymes involved in the conversion of the $\omega 3$ -fatty acid, eicosapentaenoic acid, into docosahexaenoic acid, *Biochemical Journal*, **2004**, 384, 357-366.
102. E. Sakuradani, A. Ando, J. Ogawa, S. Shimizu. Improved production of various polyunsaturated fatty acids through filamentous fungus mortierella alpina breeding, *Applied Microbiology and Biotechnology*, **2009**, 84, 1-10.
103. Q. K. Pham, H. D. Chastel. Method for mixotrophic culture of spirulinas for producing a biomass rich in omega-6 polyunsaturated fatty acids and/or in sulpholipids, United States Patent, US 0017558, 2003.
104. J. Huang, T. Akia, K. Hachida, T. Yokochi, S. Kawamoto, S. Shigeta, K. Ono, O. Suzuki.

Profile of polyunsaturated fatty acids produced by *thraustochytrium* sp. KK17-3, *Journal of the American Oil Chemists' Society*, **2001**, 78, 605-610.

105. S. M. T. Tornisielo, J. M. Vieira, M. Cecília, V. S. Carneiro, J. S. Govone. Fatty acid production by four strains of *mucor hiemalis* grown in plant oil and soluble carbohydrates, *African Journal of Biotechnology*, **2008**, 6, 1840-1847.

106. T. Aki, Y. Nagahata, K. Ishihara, Y. Tanaka, T. Morinaga, K. Higashiyama, K. Akimoto, S. Fujikawa, S. Kawamoto, S. Shigeta, K. Ono, O. Suzuki. Production of arachidonic acid by filamentous fungus, *mortierella alliacea* strain YN-15, *Journal of the American Oil Chemists' Society*, **2001**, 78, 599-604.

107. A. Nisha, G. Venkateswaran. Effect of culture variables on mycelial arachidonic acid production by *mortierella alpina*, *Food and Bioprocess Technology*, **2008**, 4, 232-240.

108. C. Ratledge, K. Kanagachandran, A. J. Anderson, D. J. Grantham, J. C. Stephenson. Production of docosahexaenoic acid by *cryptocodium cohnii* grown in a pH-auxostat culture with acetic acid as principal carbon source, *Lipids*, **2001**, 36, 1241-1246.

109. A. Mendes, P. Guerra, V. Madeira, F. Ruano, T. L. Silva, A. Reis. Study of docosahexaenoic acid production by the heterotrophic microalga *cryptocodium cohnii* CCMP 316 using carob pulp as a promising carbon source, *World Journal of Microbiology and Biotechnology*, **2007**, 23, 1209-1215.

110. H. Hoshida, T. Ohira, A. Minematsu, R. Akada, Y. Nishizawa. Accumulation of eicosapentaenoic acid in *nannochloropsis* sp. in response to elevated CO₂ concentrations, *Journal of Applied Phycology*, **2005**, 17, 29-34.

111. H. Hu, K. Gao. Response of growth and fatty acid compositions of *nannochloropsis* sp. to environmental factors under elevated CO₂ concentration, *Biotechnology Letters*, **2006**, 28, 987-992.

112. V. K. Eroshin, A. D. Satroutdinov, E. G. Dedyukhina, T. I. Chistyakova. Arachidonic acid production by *mortierella alpina* with growth-
coupled lipid synthesis, *Process Biochemistry*, **2000**, 35, 1171-1175.

113. C. Yuan, J. Wang, Y. Shang, G. Gong, J. Yao, Z. Yu. Production of arachidonic acid by *mortierella alpina* I₄₉-N₁₈, *Food Technology and Biotechnology*, **2002**, 40, 311-315.

114. B. H. Hwang, J. W. Kim, C. Y. Park, C. S. Park, Y. S. Kim, Y. W. Ryu. High-level production of arachidonic acid by fed-batch culture of *mortierella alpina* using NH₄OH as a nitrogen source and pH control, *Biotechnology Letters*, **2005**, 27, 731-735.

115. J. P. Fidalgo, A. Cid, E. Torres, A. Sukenik, C. Herrero. Effects of nitrogen source and growth phase on proximate biochemical composition, lipid classes and fatty acid profile of the marine microalga *isochrysis galbana*, *Aquaculture*, **1998**, 166, 105-116.

116. B. K. De, S. Chaudhury, D. K. Bhattacharyya. Effect of nitrogen sources on γ -linolenic acid accumulation in *spirulina platensis*, *Journal of the American Oil Chemists' Society*, **1999**, 76, 153-156.

117. K. J. Flynn, J. L. Garrido, M. Zapata, H. Opik, C. R. Hipkin. Changes in fatty acids, amino acids and carbon/nitrogen biomass during nitrogen starvation of ammonium and nitrate grown *isochrysis galbana*, *Journal of Applied Phycology*, **1992**, 4, 95-104.

118. R. Siron, G. Giusti, B. Berland. Changes in the fatty acid composition of *phaeodactylum tricornutum* and *dunaliella tertiolecta* during growth and under phosphorus deficiency, *Marine Ecology Progress Series*, **1989**, 55, 95-100.

119. I. K. Goldberg, Z. Cohen. The effect of phosphate starvation on the lipid and fatty acid composition of the freshwater eustigmatophyte *monodus subterraneus*, *Phytochemistry*, **2006**, 67, 696-701.

120. W. L. Chu, S. M. Phang, S. H. Goh. Environmental effects on growth and biochemical composition of *nitzschia inconspicua grunow*, *Journal of Applied Phycology*, **1996**, 8, 389-396.

121. L. Zhu, X. Zhang, L. Ji, X. Song, C. Kuang. Changes of lipid content and fatty acid composition of *schizochytrium limacinum* in

response to different temperatures and salinities, *Process Biochemistry*, **2007**, *42*, 210-214.

122. Y. Taoka, N. Nagano, Y. Okita, H. Izumida, S. Sugimoto, M. Hayashi. Influences of culture temperature on the growth, lipid content and fatty acid composition of *aurantiochytrium* sp. strain mh0186, *Marine Biotechnology*, **2009**, *11*, 368-374.

123. H. Jiang, K. Gao. Effects of lowering temperature during culture on the production of polyunsaturated fatty acids in the marine diatom *phaeodactylum tricornutum* (bacillariophyceae), *Journal of Phycology*, **2004**, *40*, 651-654.

124. M. A. Tedesco, E. O. Duerr. Light, temperature and nitrogen starvation effects on the total lipid and fatty acid content and composition of *spirulina platensis* UTEX 1928, *Journal of Applied Phycology*, **1989**, *1*, 201-209.

125. A. M. Nuutila, A. M. Aura, M. Kiesvaara, V. Kauppinen. The effect of salinity, nitrate concentration, pH and temperature on eicosapentaenoic acid (EPA) production by the red unicellular alga *porphyridium purpureum*, *Journal of Biotechnology*, **1997**, *55*, 55-63.

126. P. K. Bajpai, P. Bajpai, O. P. Ward. Production of arachidonic acid by *mortierella alpina* ATCC 32222, *Journal of Industrial Microbiology*, **1991**, *8*, 179-186.

127. Y. Jiang, F. Chen. Effects of medium glucose concentration and pH on docosahexaenoic acid content of heterotrophic *cryptocodonium cohnii*, *Process Biochemistry*, **2000**, *35*, 1205-1209.

128. Y. Jiang, F. Chen, S. Z. Liang. Production potential of docosahexaenoic acid by the heterotrophic marine dinoflagellate *cryptocodonium cohnii*, *Process Biochemistry*, **1999**, *34*, 633-637.

129. A. P. Carvalho, I. Pontes, H. Gaspar, F. X. Malcata. Metabolic relationships between macro- and micronutrients, and the eicosapentaenoic acid and docosahexaenoic acid contents of *pavlova lutheri*, *Enzyme and Microbial Technology*, **2006**, *38*, 358-366.

130. G. Mahajan, M. Kamat. γ -Linolenic acid production from *spirulina platensis*, *Applied Microbiology and Biotechnology*, **1995**, *43*, 466-469.

131. J. D. Weete, H. Kim, S. R. Gandhi, Y. Wang, R. Dute. Lipids and ultrastructure of *thraustochytrium* sp. ATCC 26185, *Lipids*, **1997**, *32*, 839-845.

132. A. C. Guedes, L. A. Meireles, H. M. Amaro, F. X. Malcata. Changes in lipid class and fatty acid composition of cultures of *pavlova lutheri*, in response to light intensity, *Journal of the American Oil Chemists' Society*, **2010**, *87*, 791-801.

133. I. Tzovenis, N. D. Pauw, P. Sorgeloos. Optimisation of T-ISO biomass production rich in essential fatty acids II, effect of different light regimes on the production of fatty acids, *Aquaculture*, **2003**, *216*, 223-242.

134. Z. Xu, X. Yan, L. Pei, Q. Luo, J. Xu. Changes in fatty acids and sterols during batch growth of *pavlova viridis* in photobioreactor, *Journal of Applied Phycology*, **2008**, *20*, 237-243.

135. R. B. Bailey, D. DiMasi, J. M. Hansen, P. J. Mirrasoul, C. M. Ruecker, G. T. Veeder, T. Kaneko, W. R. Barclay. Enhanced production of lipids containing polyenoic fatty acid by very high density cultures of eukaryotic microbes in fermenters, United States Patent, US 8216812B2, 2006.

136. R. Ruenwai, S. Cheevadhanarak, S. Rachdawong, M. Tanticharoen, K. Laoeng. Oxygen-induced expression of Δ^6 -, Δ^9 - and Δ^{12} -desaturase genes modulates fatty acid composition in *mucor rouxii*, *Applied Microbiology and Biotechnology*, **2010**, *86*, 327-334.

137. S. Y. Ho, F. Chen. Lipid characterization of *mortierella alpina* grown at different NaCl concentrations, *Journal of Agricultural and Food Chemistry*, **2008**, *56*, 7903-7909.

138. G. Q. Chen, Y. Jiang, F. Chen. Fatty acid and lipid class composition of the eicosapentaenoic acid-producing microalga *nitzschia laevis*, *Food Chemistry*, **2007**, *104*, 1580-1585.

139. G. Q. Chen, Y. Jiang, F. Chen. Salt-induced alterations in lipid composition of diatom *nitzschia laevis* (bacillariophyceae) under heterotrophic culture condition, *Journal of Phycology*, **2008**, *44*, 1309-1314.

140. G. Q. Chen, Y. Jiang, F. Chen. Variation of lipid class composition in *nitzschia laevis* as a response to growth temperature change, *Food Chemistry*, **2008**, *109*, 88-94.

141. R. J. Henderson, J. W. Leftley, J. R. Sargent. Lipid composition and biosynthesis in the marine dinoflagellate *cryptocodinium cohnii*, *Phytochemistry*, **1988**, *27*, 1679-1683.

142. Z. Cohen, M. Reungjitchachawali, W. Siangdung, M. Tantcharoen. Production and partial purification of γ -linolenic acid and some pigments from *spirulina platensis*, *Journal of Applied Phycology*, **1993**, *5*, 109-115.

143. Y. Kamisaka, N. Noda, T. Sakai, K. Kawasaki. Lipid bodies and lipid body formation in an oleaginous fungus, *mortierella ramanniana* var. *angulispora*, *Biochimica et Biophysica Acta*, **1999**, *1438*, 185-198.

144. E. Morita, Y. Kumon, T. Nakahara, S. Kagiwada, T. Noguchi. Docosahexaenoic acid production and lipid-body formation in *schizochytrium limacinum* SR21, *Marine Biotechnology*, **2006**, *38*, 319-327.

145. C. P. Liu, L. P. Lin. Ultrastructural study and lipid formation of *isochrysis* sp. CCMP1324, *Botanical Bulletin of Academia Sinica*, **2001**, *42*, 207-214.

146. M. J. Jin, H. Huang, A. H. Xiao, Z. Gao, X. Liu, C. Peng. Enhancing arachidonic acid production by *nortierella alpina* ME-1 using improved mycelium aging technology, *Bioprocess Biosyst Eng*, **2009**, *32*, 117-122.

147. Z. C. Wu, Z. Cohen, I. K. Goldberg, A. Richmond. Characterization of growth and arachidonic acid production of *parietochloris incisa* comb. nov (trebouxiophyceae, chlorophyta), *Journal of Applied Phycology*, **2002**, *14*, 453-460.

148. S. Shimizu, H. Kawashima, Y. Shinmen, K. Akimoto, H. Yamada. Microbial conversion of an oil containing γ -linolenic acid to an oil containing eicosapentaenoic acid, *Journal of the American Oil Chemists' Society*, **1989**, *66*, 342-347.

149. J. M. F. Sevilla, M. C. C. García, A. S. Mirón, E. H. Belarbi, F. G. Camacho, E. M. Grima. Pilot-plant-scale outdoor mixotrophic cultures of *phaeodactylum tricornutum* using glycerol in vertical bubble column and airlift photobioreactors: studies in fed-batch mode, *Biotechnology Progress*, **2004**, *20*, 728-736.

150. J. G. Burgess, K. Iwamoto, Y. Miura, H. Takano, T. Matsunaga. An optical fibre photobioreactor for enhanced production of the marine unicellular alga *isochrysis* aft. *galbana* T-Iso (UTEX LB 2307) rich in docosahexaenoic acid, *Applied Microbiology and Biotechnology*, **1993**, *39*, 456-459.

151. T. E. Lewis, P. D. Nichols, T. A. McMeekin. The biotechnological potential of *thraustochytrids*, *Marine Biotechnology*, **1999**, *1*, 580-587.

152. F. Xu, Z. L. Cai, W. Cong, F. Ouyang. Growth and fatty acid composition of *nannochloropsis* sp. grown mixotrophically in fed-batch culture, *Biotechnology Letters*, **2004**, *26*, 1319-1322.

153. M. Akimoto, A. Shirai, K. Ohtaguchi, K. Koide. Carbon dioxide fixation and polyunsaturated fatty acid production by the red alga *porphyridium cruentum*, *Applied Biochemistry and Biotechnology*, **1998**, *73*, 269-278.

154. F. G. A. Fernández, F. G. Camacho, J. A. S. Pérez, J. M. F. Sevilla, E. M. Grima. Modeling of eicosapentaenoic acid (EPA) production from *phaeodactylum tricornutum* cultures in tubular photobioreactors, effects of dilution rate, tube diameter and solar irradiance, *Biotechnology & Bioengineering*, **2000**, *68*, 173-183.

155. E. M. Grima, J. A. S. Pérez, F. G. Camacho, J. U. Cardona, J. M. F. Sevilla, F. G. A. Fernández. Biomass and eicosapentaenoic acid productivities from an outdoor batch culture of *phaeodactylum tricornutum* UTEX 640 in an airlift tubular photobioreactor, *Applied Microbiology and Biotechnology*, **1995**, *42*, 658-663.

156. A. Meiser, U. S. Staiger, W. Trosch. Optimization of eicosapentaenoic acid production by *phaeodactylum tricornutum* in the flat panel airlift (FPA) reactor, *Journal of Applied Phycology*, **2004**, *16*, 215-225.

157. E. M. Grima, F. G. G. Camacho, J. A. S. Pérez, J. U. Cardona, F. G. A. Fernández, J. M. F. Sevilla. Outdoor chemostat culture of *phaeodactylum tricornutum* UTEX 640 in a tubular photobioreactor for the production of eicosapentaenoic acid, *Biotechnology and Applied Biochemistry*, **1994**, 20, 279-290.

158. A. Reis, L. Gouveia, V. Veloso, H. L. Fernandes, J. A. Empis, J. M. Novais. Eicosapentaenoic acid-rich biomass production by the microalga *phaeodactylum tricornutum* in a continuous-flow reactor, *Bioresource Technology*, **1996**, 55, 83-88.

159. A. P. Carvalho, F. X. Malcata. Effect of culture media on production of polyunsaturated fatty acids by *pavlova lutheri*, *Cryptogamie Algologie*, **2000**, 21, 59-71.

160. E. M. Grima, J. A. S. Pérez, J. L. G. Sánchez, F. G. Camacho, D. L. Alonso. EPA from *isochrysis galbana*, growth conditions and productivity, *Process Biochemistry*, **1992**, 27, 299-305.

161. E. M. Grima, J. A. S. Pérez, F. G. Camacho, J. M. F. Sevilla, F. G. A. Fernández. Effect of growth rate on the eicosapentaenoic acid and docosahexaenoic acid content of *isochrysis galbana* in chemostat culture, *Applied Microbiology and Biotechnology*, **1994**, 41, 23-27.

162. E. M. Grima, J. A. S. Pérez, F. G. Camacho, J. L. G. Sánchez, D. L. Alonso. N-3 PUFA productivity in chemostat cultures of microalgae, *Applied Microbiology and Biotechnology*, **1993**, 38, 599-605.

163. Y. H. Lin, F. L. Chang, C. Y. J. Y. Tsao, Leu. Influence of growth phase and nutrient source on fatty acid composition of *isochrysis galbana* CCMP 1324 in a batch photoreactor, *Biochemical Engineering Journal*, **2007**, 37, 166-176.

164. L. Poisson, F. Ergan. Docosahexaenoic acid ethyl esters from *isochrysis galbana*, *Journal of Biotechnology*, **2001**, 91, 75-81.

165. E. Gánuza, M. S. Izquierdo. Lipid accumulation in *schizochytrium G13/2S* produced in continuous culture, *Applied Microbiology and Biotechnology*, **2007**, 76, 985-990.

166. R. D. Bowles, A. E. Hunt, G. B. Bremer, M. G. Duchars, R. A. Eaton. Long-chain n-3 polyunsaturated fatty acid production by members of the marine protistan group the thraustochytrids: screening of isolates and optimisation of docosahexaenoic acid production, *Journal of Biotechnology*, **1999**, 70, 193-202.

167. A. P. Carvalho, S. O. Silva, J. M. Baptista, F. X. Malcata. Light requirements in microalgal photobioreactors: an overview of biophotonic aspects, *Applied Microbiology and Biotechnology*, **2011**, 89, 1275-1288.

168. A. Singh, O. P. Ward. Production of high yields of arachidonic acid in a fed-batch system by *mortierella alpina* ATCC 32222, *Applied Microbiology and Biotechnology*, **1997**, 48, 1-5.

169. Y. Zeng, X. J. Ji, M. Lian, L. J. Ren, L. J. Jin, P. K. Ouyang, H. Huang. Development of a temperature shift strategy for efficient docosahexaenoic acid production by a marine fungoid protist, *schizochytrium* sp. HX-308, *Applied Biochemistry and Biotechnology*, **2011**, 164, 249-255.