

# **Phép chiếu Bergman tác động lên $L^\infty$ trên hình cầu đơn vị $\mathbb{B}_n$**

**Lê Văn An\***

*Khoa Toán và Thống kê, Trường Đại học Quy Nhơn, Việt Nam.*

*Ngày nhận bài: 31/10/2021, Ngày nhận đăng: 24/12/2021*

## **TÓM TẮT**

Với một hàm trọng cho trước, ta định nghĩa phép chiếu kiểu Bergman cảm sinh bởi hàm trọng đó. Chúng tôi đưa ra một đặc trưng của hàm trọng bán kính để phép chiếu kiểu Bergman đó là bị chặn từ  $L^\infty$  vào không gian Bloch  $\mathcal{B}$  trên hình cầu đơn vị  $\mathbb{B}_n$  của  $\mathbb{C}^n, n > 1$ .

**Từ khóa:** *Không gian Bergman, phép chiếu Bergman, không gian Bloch.*

---

\*Tác giả liên hệ chính.

Email: [levanan@qnu.edu.vn](mailto:levanan@qnu.edu.vn)

# On the Bergman projections acting on $L^\infty$ on the unit ball $\mathbb{B}_n$

Van An Le\*

Department of Mathematics and Statistics, Quy Nhon University, Vietnam.

Received: 31/10/2021; Accepted: 24/12/2021

## ABSTRACT

Given a weight function, we define the Bergman type projection induced by this weight. We characterize the radial weights such that this projection is bounded from  $L^\infty$  to the Bloch space  $\mathcal{B}$  on the unit ball  $\mathbb{B}_n$  of  $\mathbb{C}^n$ ,  $n > 1$ .

**Keywords:** Bergman space, Bergman projection, Bloch space.

## 1. INTRODUCTION AND MAIN RESULT

Let  $\mathbb{C}^n$  denote the  $n$ -dimensional complex Euclidean space. For any two points  $z = (z_1, \dots, z_n)$ ,  $w = (w_1, \dots, w_n)$  in  $\mathbb{C}^n$ , we use the well-known notation

$$\langle z, w \rangle = z_1 \overline{w_1} + \dots + z_n \overline{w_n},$$

and  $|z| = \sqrt{\langle z, z \rangle}$ .

Let  $\mathbb{B}_n = \{z \in \mathbb{C}^n : |z| < 1\}$  be the unit ball, and let  $\mathbb{S}_n = \{z \in \mathbb{C}^n : |z| = 1\}$  be the unit sphere in  $\mathbb{C}^n$ .

Denote by  $H(\mathbb{B}_n)$  the space of all holomorphic functions on the unit ball  $\mathbb{B}_n$ . Let  $dv$  be the normalized volume measure on  $\mathbb{B}_n$ . The normalized surface measure on  $\mathbb{S}_n$  will be denoted by  $d\sigma$ .

Let  $\rho$  be a positive and integrable function on  $[0, 1)$ . We extend it to  $\mathbb{B}_n$  by  $\rho(z) = \rho(|z|)$ , and call such  $\rho$  a radial weight function. The weighted Bergman space  $A_\rho^2$  is the space of functions  $f$  in  $H(\mathbb{B}_n)$  such that

$$\|f\|_\rho^2 = \int_{\mathbb{B}_n} |f(z)|^2 \rho(z) dv(z) < \infty.$$

Let  $\rho$  be a radial weight and  $X$  be a space of measurable functions on  $\mathbb{B}_n$ . The Bergman type projection  $P_\rho$  acting on  $X$  is given by

$$P_\rho f(z) = \int_{\mathbb{B}_n} K_\rho(z, w) f(w) \rho(w) dv(w),$$

for  $z \in \mathbb{B}_n$ ,  $f \in X$ , where  $K_\rho(z, w)$  is the reproducing kernel of the weighted Bergman space  $A_\rho^2$ .

When  $\rho$  is the standard radial weight  $\rho(z) = (1 - |z|^2)^\alpha$ ,  $\alpha > -1$ , the corresponding projection is denoted by  $P_\alpha$ .

Corresponding author.

Email: levanan@qnu.edu.vn

A radial weight  $\rho$  belongs to the class  $\widehat{\mathcal{D}}$  if  $\widehat{\rho}(r) \lesssim \widehat{\rho}(\frac{1+r}{2})$  for all  $r \in [0, 1)$ , where  $\widehat{\rho}(r) = \int_r^1 \rho(s)ds$ .

The Bergman space and a lot of topics related to this famous space have attracted the attention of many mathematicians so far. When the weight function is the standard radial weight, the theory of Bergman space is well-known. We recommend the books of Zhu<sup>9,10</sup> for more details.

In 2018, we began the study of small Bergman spaces in higher dimensions<sup>5</sup>. The projections play a crucial role in studying operator theory on spaces of analytic functions. Bounded analytic projections can also be used to establish duality relations and to obtain useful equivalent norms in spaces of analytic functions. Hence the boundedness of projections is an interesting topic which has been studied by many authors in recent years<sup>1,2,3,7,8</sup>. In<sup>7</sup>, Peláez and Rättyä considered the projection  $P_{\rho_1}$  acting on  $L_{\rho_2}^p(\mathbb{D})$ ,  $1 \leq p < \infty$ , when two weights  $\rho_1, \rho_2$  are in the class  $\mathcal{R}$  of the so called regular weights. A radial weight  $\rho$  is regular if  $\widehat{\rho}(r) \asymp (1-r)\rho(r)$ ,  $r \in (0, 1)$ . Recently, in 2019, they extended these results to the case where  $\rho_1 \in \widehat{\mathcal{D}}$ ,  $\rho_2$  is radial<sup>8</sup>.

In this paper, we are going to study the projections acting on the space  $L^\infty$ . Let us recall that the Bloch space of  $\mathbb{B}_n$ , denoted by  $\mathcal{B}(\mathbb{B}_n)$ , or simply by  $\mathcal{B}$ , is the space of holomorphic functions  $f$  on  $\mathbb{B}_n$  such that

$$\sup_{z \in \mathbb{B}_n} (1-|z|^2) |Rf(z)| < \infty,$$

where

$$Rf(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z)$$

is the radial derivative of  $f$  at  $z \in \mathbb{B}_n$ . In the one dimensional case, the Bloch space consists of analytic functions  $f$  on  $\mathbb{D}$  such that

$$\sup_{z \in \mathbb{D}} (1-|z|^2) |f'(z)| < \infty,$$

and is denoted by  $\mathcal{B}(\mathbb{D})$ .

In the case of standard radial weight, we have the following result.

**Theorem A.** *For any  $\alpha > -1$ , the Bergman type projection  $P_\alpha$  is a bounded linear operator from  $L^\infty$  onto the Bloch space  $\mathcal{B}$ .*

This theorem is also valid for the case of one dimension  $\mathbb{C}$  and of higher dimension  $\mathbb{C}^n$ . See<sup>10</sup> Theorem 5.2 for the proof in the case of one variable and<sup>9</sup> Theorem 3.4 for the proof in the case of several variables.

When the weight functions is more general, in<sup>8</sup>, Peláez and Rättyä obtained an interesting result in the one dimensional case.

**Theorem B.** *Let  $\rho$  be a radial weight. Then the projection  $P_\rho : L^\infty(\mathbb{D}) \rightarrow \mathcal{B}(\mathbb{D})$  is bounded if and only if  $\rho \in \widehat{\mathcal{D}}$ .*

The aim of this paper is to study the Bergman type projections acting on  $L^\infty$  in the case of higher dimension. In this case, what will be the target space and the characterizations of the weight functions such that the projection is bounded? We extend Theorem B to the case of several variables and obtain the following result.

**Theorem 1.1.** *Let  $\rho$  be a radial weight. Then the projection  $P_\rho : L^\infty \rightarrow \mathcal{B}$  is bounded if and only if  $\rho \in \widehat{\mathcal{D}}$ .*

The paper is organized as follows. In Section 2, we give some important lemmas which are used in the proof of Theorem 1.1 and Section 3 gives us an explicit explanation for our main result.

Throughout this text, the notation  $U(z) \lesssim V(z)$  (or equivalently  $V(z) \gtrsim U(z)$ ) means that there is a positive constant  $C$  such that  $U(z) \leq C.V(z)$  holds for all  $z$  in

the set in question, which may be a space of functions or a set of numbers. If both  $U(z) \lesssim V(z)$  and  $V(z) \lesssim U(z)$  hold, then we write  $U(z) \asymp V(z)$ .

## 2. SOME AUXILIARY LEMMAS

To prove Theorem 1.1 we need several auxiliary lemmas.

**Lemma 2.1.** *Let  $\rho$  be a radial weight. Then the following conditions are equivalent:*

- (i)  $\rho \in \widehat{\mathcal{D}}$ ;
- (ii) *There exist  $C = C(\rho) > 0$  and  $\beta_0 = \beta_0(\rho) > 0$  such that*

$$\widehat{\rho}(r) \leq C \left( \frac{1-r}{1-t} \right)^\beta \widehat{\rho}(t),$$

*where  $0 \leq r \leq t < 1$ , for all  $\beta \geq \beta_0$ ;*

- (iii) *The asymptotic equality*

$$\int_0^1 s^x \rho(s) ds \asymp \widehat{\rho} \left( 1 - \frac{1}{x} \right),$$

*where  $x \in [1, \infty)$ , is valid;*

- (iv) *There exist  $C_0 = C_0(\rho) > 0$  and  $C = C(\rho) > 0$  such that*

$$\widehat{\rho}(0) \leq C_0 \widehat{\rho} \left( \frac{1}{2} \right)$$

*and  $\rho_n \leq C \rho_{2n}$  for all  $n \in \mathbb{N}$ .*

Lemma 2.1 gives us the characterizations of the weight function  $\rho \in \widehat{\mathcal{D}}$ . The proof of this lemma can be found in<sup>6</sup>.

**Lemma 2.2.** *If*

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \in H^p, \quad 0 < p \leq 2,$$

*where  $H^p$  denotes the Hardy space, then*

$$\sum_{j=0}^{\infty} (j+1)^{p-2} |a_j|^p \lesssim \|f\|_p^p.$$

**Lemma 2.3.** *Let  $\{a_j\}$  be a sequence of complex numbers such that  $\sum j^{q-2} |a_j|^q < \infty$  for some  $q, 2 \leq q < \infty$ . Then the function  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  is in  $H^q$ , and*

$$\|f\|_q^q \lesssim \sum_{j=0}^{\infty} (j+1)^{q-2} |a_j|^q.$$

Two above lemmas are the classical Hardy-Littlewood inequalities, which can be found, for example, in Duren's book<sup>4</sup> Theorem 6.2 and 6.3.

In the following lemma, we give the explicit formula for the reproducing kernel of  $A_\rho^2$ .

**Lemma 2.4.** *Let  $\rho$  be a radial weight. Then the reproducing kernel  $K_\rho(z, w)$  is given by*

$$K_\rho(z, w) = \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d! n! \rho_{2n-1+2d}} \langle z, w \rangle^d,$$

*for  $z, w \in \mathbb{B}_n$ , where*

$$\rho_x = \int_0^1 t^x \rho(t) dt, \quad x \geq 1.$$

*Proof.* By the multinomial formula (see<sup>9</sup> (1.1)), we have that

$$\langle z, w \rangle^d = \sum_{\beta \in \mathbb{N}^n, |\beta|=d} \frac{d!}{\beta!} z^\beta \bar{w}^\beta, \quad z, w \in \mathbb{C}^n,$$

where  $|\beta| = \beta_1 + \dots + \beta_n$ ,  $\beta! = \beta_1! \dots \beta_n!$  for each  $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{N}^n$ . Hence, for  $\alpha \in \mathbb{N}^n$ ,  $|\alpha| = d$ ,

$$\begin{aligned} \int_{\mathbb{S}_n} \xi^\alpha \langle z, \xi \rangle^d d\sigma(\xi) \\ = \sum_{\beta \in \mathbb{N}^n, |\beta|=d} \frac{d! z^\beta}{\beta!} \int_{\mathbb{S}_n} \xi^\alpha \bar{\xi}^\beta d\sigma(\xi), \end{aligned}$$

where  $z \in \mathbb{B}_n$ . By Lemma 1.11 in<sup>9</sup>,

$$\int_{\mathbb{S}_n} \xi^\alpha \bar{\xi}^\beta d\sigma(\xi) = \begin{cases} 0 & \text{if } \alpha \neq \beta, \\ \frac{\alpha!(n-1)!}{(d+n-1)!} & \text{if } \alpha = \beta, \end{cases}$$

and we obtain

$$\begin{aligned} \int_{\mathbb{S}_n} \xi^\alpha \langle z, \xi \rangle^d d\sigma(\xi) &= \frac{d!}{\alpha!} z^\alpha \int_{\mathbb{S}_n} \xi^\alpha \bar{\xi}^\alpha d\sigma(\xi) \\ &= \frac{d!}{\alpha!} \frac{\alpha!(n-1)!}{(d+n-1)!} z^\alpha \\ &= \frac{d!(n-1)!}{(d+n-1)!} z^\alpha, \end{aligned}$$

for  $z \in \mathbb{B}_n$ . Therefore, for  $\alpha \in \mathbb{N}^n$ ,  $|\alpha| = d$  we have

$$\begin{aligned} &\int_{\mathbb{B}_n} w^\alpha \langle z, w \rangle^d \rho(w) dv(w) \\ &= 2n \int_0^1 t^{2n-1+2d} \rho(t) dt \int_{\mathbb{S}_n} \xi^\alpha \langle z, \xi \rangle^d d\sigma(\xi) \\ &= \frac{2d!n!\rho_{2n-1+2d}}{(d+n-1)!} z^\alpha, \quad z \in \mathbb{B}_n. \end{aligned}$$

It follows that

$$z^\alpha = \frac{(d+n-1)!}{2d!n!\rho_{2n-1+2d}} \int_{\mathbb{B}_n} w^\alpha \langle z, w \rangle^d \rho(w) dv(w), \quad (1)$$

for any  $z \in \mathbb{B}_n$ .

Since  $\rho(t) > 0$ ,  $0 < t < 1$ , for every  $\varepsilon > 0$  there exists  $C_\varepsilon > 0$  such that  $\rho_s \geq C_\varepsilon(1-\varepsilon)^s$ . Given  $z \in \mathbb{B}_n$ , we have

$$\begin{aligned} &\int_{\mathbb{B}_n} \left| \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d!n!\rho_{2n-1+2d}} \langle z, w \rangle^d \right|^2 \rho(w) dv(w) \\ &= \frac{1}{4} \sum_{d_1, d_2 \geq 0} \frac{(d_1+n-1)!(d_2+n-1)!}{d_1!d_2!(n!)^2 \rho_{2n-1+2d_1} \rho_{2n-1+2d_2}} \times \\ &\quad \times \int_{\mathbb{B}_n} \langle z, w \rangle^{d_1} \langle w, z \rangle^{d_2} \rho(w) dv(w) \\ &= \frac{1}{4} \sum_{d_1, d_2 \geq 0} \frac{(d_1+n-1)!(d_2+n-1)!}{d_1!d_2!(n!)^2 \rho_{2n-1+2d_1} \rho_{2n-1+2d_2}} \times \\ &\quad \times \int_{\mathbb{B}_n} \sum_{|\beta|=d_2} w^\beta \bar{z}^\beta \frac{d_2!}{\beta!} \langle z, w \rangle^{d_1} \rho(w) dv(w) \\ &= \frac{1}{2} \sum_{d \geq 0} \left( \frac{(d+n-1)!}{d!n!} \right) \frac{1}{\rho_{2n-1+2d}^2} \times \end{aligned}$$

$$\begin{aligned} &\times \sum_{|\beta|=d} \frac{(d!)^2}{\beta!} \frac{n!\rho_{2n-1+2d}}{(d+n-1)!} z^\beta \bar{z}^\beta \\ &= \frac{1}{2} \sum_{d \geq 0} \frac{(d+n-1)!}{n!\rho_{2n-1+2d}} \sum_{|\beta|=d} \frac{z^\beta \bar{z}^\beta}{\beta!} \\ &= \frac{1}{2} \sum_{d \geq 0} \frac{(d+n-1)!}{d!n!\rho_{2n-1+2d}} |z|^{2d} < \infty. \end{aligned}$$

Thus, the function

$$w \mapsto \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d!n!\rho_{2n-1+2d}} \langle w, z \rangle^d$$

belongs to  $A_\rho^2$ .

By (1) and by continuity, for every  $f \in A_\rho^2(\mathbb{B}_n)$ , we have

$$\begin{aligned} f(z) &= \int_{\mathbb{B}_n} f(w) \times \\ &\times \left( \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d!n!\rho_{2n-1+2d}} \langle z, w \rangle^d \right) \rho(w) dv(w), \end{aligned}$$

for  $z \in \mathbb{B}_n$ .

Therefore,

$$K_\rho(z, w) = \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d!n!\rho_{2n-1+2d}} \langle z, w \rangle^d.$$

□

### 3. PROOF OF MAIN RESULT

It suffices to consider only the case  $n > 1$ .

**Proposition 3.1.** *If  $\rho \in \widehat{\mathcal{D}}$ , then the projection  $P_\rho : L^\infty \rightarrow \mathcal{B}$  is bounded, where  $P_\rho$  is defined by*

$$P_\rho \varphi(z) = \int_{\mathbb{B}_n} K_\rho(z, w) \varphi(w) \rho(w) dv(w),$$

for  $\varphi \in L^\infty$ ,  $z \in \mathbb{B}_n$ .

*Proof.* We have

$$K_\rho(z, w) = \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d! n! \rho_{2n-1+2d}} \langle z, w \rangle^d.$$

Hence, for a fixed  $w \in \mathbb{B}_n$ ,

$$\begin{aligned} & RK_\rho(z, w) \\ &= \sum_{j=1}^n z_j \frac{\partial K_\rho(z, w)}{\partial z_j} \\ &= \sum_{j=1}^n z_j \frac{\partial}{\partial z_j} \left( \frac{1}{2} \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d! n! \rho_{2n-1+2d}} \langle z, w \rangle^d \right) \\ &= \frac{1}{2} \sum_{j=1}^n z_j \sum_{d=0}^{\infty} \frac{(d+n-1)!}{d! n! \rho_{2n-1+2d}} d\bar{w}_j \langle z, w \rangle^{d-1} \\ &= \frac{1}{2} \sum_{d=1}^{\infty} \frac{(d+n-1)!}{(d-1)! n! \rho_{2n-1+2d}} \langle z, w \rangle^d \\ &= \frac{1}{2} \sum_{d=1}^{\infty} \frac{\Gamma(d+n)}{\Gamma(d)\Gamma(n+1)\rho_{2n-1+2d}} \langle z, w \rangle^d. \end{aligned}$$

Now, given  $\varphi \in L^\infty$ , let

$$\begin{aligned} f(z) &:= P_\rho \varphi(z) \\ &= \int_{\mathbb{B}_n} K_\rho(z, w) \varphi(w) \rho(w) dv(w), \end{aligned}$$

where  $z \in \mathbb{B}_n$ . For all  $z \in \mathbb{B}_n$  we have

$$\begin{aligned} |Rf(z)| &= \left| \int_{\mathbb{B}_n} RK_\rho(z, w) \varphi(w) \rho(w) dv(w) \right| \\ &\leq \int_{\mathbb{B}_n} |RK_\rho(z, w)| |\varphi(w)| \rho(w) dv(w) \\ &\leq \|\varphi\|_\infty \int_{\mathbb{B}_n} |RK_\rho(z, w)| \rho(w) dv(w). \end{aligned} \tag{2}$$

Set

$$g(\lambda) = \sum_{d=1}^{\infty} \frac{\Gamma(d+n)}{\Gamma(d)} \frac{\lambda^{d-1}}{\rho_{2n-1+2d}}, \quad \lambda \in \mathbb{D}.$$

Since  $\rho(t) > 0, 0 < t < 1$ ,  $g$  is analytic on the unit disk. Then

$$RK_\rho(z, w) = \frac{\langle z, w \rangle}{2\Gamma(n+1)} g(\langle z, w \rangle). \tag{3}$$

Next we consider the reproducing kernel  $K_\rho^1(z, w)$  of the Bergman space in the unit disk with the weight  $\rho$ . We have

$$K_\rho^1(z, w) = \frac{1}{2} \sum_{d=0}^{\infty} \frac{(z\bar{w})^d}{\rho_{2d+1}}.$$

Furthermore,

$$\begin{aligned} \frac{\partial^n}{\partial z^n} K_\rho^1(z, w) &= \frac{1}{2} \sum_{d=n}^{\infty} \frac{\Gamma(d+1)(z\bar{w})^{d-n}\bar{w}^n}{\Gamma(d-n+1)\rho_{2d+1}} \\ &= \frac{1}{2} \sum_{s=1}^{\infty} \frac{\Gamma(s+1)}{\Gamma(s)} \frac{(z\bar{w})^{s-1}\bar{w}^n}{\rho_{2s+2n-1}} \\ &= \frac{1}{2} g(z\bar{w})\bar{w}^n. \end{aligned}$$

By a result of Peláez and Rättyä (<sup>7</sup> Theorem 1 (ii)), we have

$$\begin{aligned} \int_{\mathbb{D}} \left| \frac{\partial^n}{\partial z^n} K_\rho^1(z, w) \right| (1-|z|^2)^{n-2} dA(z) \\ \asymp \int_0^{|w|} \frac{dt}{\widehat{\rho}(t)(1-t)^2}, \quad \frac{1}{2} \leq |w| < 1, \end{aligned}$$

where  $\widehat{\rho}(t) = \int_t^1 \rho(s) ds$ .

Thus,

$$\begin{aligned} \int_{\mathbb{D}} |g(z\bar{w})| (1-|z|^2)^{n-2} dA(z) \\ \asymp \int_0^{|w|} \frac{dt}{\widehat{\rho}(t)(1-t)^2}, \quad \frac{1}{2} \leq |w| < 1. \end{aligned}$$

Since  $g$  is analytic on the unit disk, we have

$$\begin{aligned} \int_{\mathbb{D}} |g(z\bar{w})| (1-|z|^2)^{n-2} dA(z) \\ \lesssim 1 + \int_0^{|w|} \frac{dt}{\widehat{\rho}(t)(1-t)^2}, \quad w \in \mathbb{D}. \tag{4} \end{aligned}$$

Now, by (3), we have

$$\int_{\mathbb{B}_n} |RK_\rho(z, w)| \rho(w) dv(w)$$

$$\begin{aligned} &\lesssim \int_{\mathbb{B}_n} |g(\langle z, w \rangle)| \rho(w) dv(w) \\ &\asymp \int_0^1 r^{2n-1} \rho(r) \left( \int_{\mathbb{S}_n} |g(\langle rz, \xi \rangle)| d\sigma(\xi) \right) dr. \end{aligned}$$

By<sup>9</sup> Lemma 1.9 and the unitary invariance of  $d\sigma$ , we have

$$\begin{aligned} &\int_{\mathbb{S}_n} |g(\langle rz, \xi \rangle)| d\sigma(\xi) \\ &\asymp \int_{\mathbb{D}} |g(r|z| \lambda)| (1 - |\lambda|^2)^{n-2} dA(\lambda). \end{aligned}$$

Thus, by (4) we obtain

$$\begin{aligned} &\int_{\mathbb{B}_n} |RK_\rho(z, w)| \rho(w) dv(w) \\ &\lesssim \int_0^1 r^{2n-1} \rho(r) \left( 1 + \int_0^{r|z|} \frac{dt}{\hat{\rho}(t)(1-t)^2} \right) dr \\ &\lesssim 1 + \int_0^{|z|} \frac{1}{\hat{\rho}(t)(1-t)^2} \left( \int_{t \wedge |z|}^1 r^{2n-1} \rho(r) dr \right) dt \\ &\lesssim 1 + \int_0^{|z|} \frac{\hat{\rho}(t/|z|)}{\hat{\rho}(t)} \frac{dt}{(1-t)^2} \\ &\lesssim \frac{1}{1-|z|}, \quad z \in \mathbb{B}_n. \end{aligned}$$

By (2) we obtain now that

$$|Rf(z)| \lesssim \|\varphi\|_\infty \frac{1}{1-|z|^2}, \quad z \in \mathbb{B}_n.$$

Hence,

$$\sup_{z \in \mathbb{B}_n} (1-|z|^2) |Rf(z)| \lesssim \|\varphi\|_\infty.$$

It is easy to see that

$$|f(0)| \lesssim \|\varphi\|_\infty.$$

Therefore,  $P_\rho$  is bounded. The Proposition 3.1 is proved.  $\square$

**Proposition 3.2.** *Suppose that the projection  $P_\rho : L^\infty \rightarrow \mathcal{B}$  is bounded. Then  $\rho \in \widehat{\mathcal{D}}$ .*

*Proof.* Given  $\xi \in \mathbb{S}_n$  and  $w \in \mathbb{B}_n$ , let us consider a function  $g$  given by

$$g(\lambda) = RK_\rho(\lambda \xi, w), \quad \lambda \in \mathbb{D}.$$

Then

$$g(\lambda) = \sum_{d=1}^{\infty} c_d \langle \xi, w \rangle^d \lambda^d,$$

$$\text{where } c_d = \frac{1}{2n} \frac{\Gamma(d+n)}{\Gamma(d)\Gamma(n)\rho_{2n-1+2d}}.$$

By the Hardy–Littlewood inequality (see Lemma 2.2) we have

$$\begin{aligned} \sum_{d=1}^{\infty} \frac{c_d |\langle \xi, w \rangle|^d}{d+1} &\lesssim \int_0^{2\pi} |g(e^{i\theta})| \frac{d\theta}{2\pi} \\ &= \int_0^{2\pi} |RK_\rho(e^{i\theta} \xi, w)| \frac{d\theta}{2\pi}. \end{aligned}$$

Integrating both sides of the above inequality over  $\xi \in \mathbb{S}_n$  we obtain

$$\begin{aligned} &\sum_{d=1}^{\infty} \frac{c_d}{d+1} \int_{\mathbb{S}_n} |\langle \xi, w \rangle|^d d\sigma(\xi) \\ &\lesssim \int_{\mathbb{S}_n} \int_0^{2\pi} |RK_\rho(e^{i\theta} \xi, w)| \frac{d\theta}{2\pi} d\sigma(\xi) \\ &= \int_{\mathbb{S}_n} |RK_\rho(\xi, w)| d\sigma(\xi). \end{aligned}$$

By the unitary invariance of  $d\sigma$  and<sup>9</sup> Lemma 1.9, we have

$$\begin{aligned} &\int_{\mathbb{S}_n} |\langle \xi, w \rangle|^d d\sigma(\xi) \\ &= |w|^d \int_{\mathbb{S}_n} |\xi_1|^d d\sigma(\xi) \\ &= (n-1) |w|^d \int_{\mathbb{D}} (1-|z|^2)^{n-2} |z|^d dA(z) \\ &= (n-1) \pi |w|^d \int_0^1 (1-t)^{n-2} t^{d/2} dt \\ &\asymp \frac{\Gamma(\frac{d}{2}+1)\Gamma(n)}{\Gamma(\frac{d}{2}+n)} |w|^d. \end{aligned}$$

Hence,

$$\begin{aligned} &\int_{\mathbb{S}_n} |RK_\rho(\xi, w)| d\sigma(\xi) \\ &\gtrsim \sum_{d=1}^{\infty} \frac{c_d}{d+1} \frac{\Gamma(\frac{d}{2}+1)\Gamma(n)}{\Gamma(\frac{d}{2}+n)} |w|^d \end{aligned}$$

$$= \frac{1}{2n} \sum_{d=1}^{\infty} \frac{\Gamma(d+n)\Gamma(\frac{d}{2}+1)}{(d+1)\Gamma(d)\Gamma(\frac{d}{2}+n)\rho_{2n-1+2d}} |w|^d.$$

Since

$$\frac{\Gamma(d+n)\Gamma(\frac{d}{2}+1)}{(d+1)\Gamma(d)\Gamma(\frac{d}{2}+n)} \asymp 1,$$

we get

$$\begin{aligned} \int_{\mathbb{S}_n} |RK_{\rho}(\xi, w)| d\sigma(\xi) \\ \gtrsim \frac{1}{2n} \sum_{d=1}^{\infty} \frac{|w|^d}{\rho_{2n-1+2d}}, \quad w \in \mathbb{B}_n. \end{aligned}$$

Therefore, for  $z \in \mathbb{B}_n$ , we have

$$\begin{aligned} \int_{\mathbb{B}_n} |RK_{\rho}(z, w)| \rho(w) dv(w) \\ = 2n \int_0^1 r^{2n-1} \rho(r) \int_{\mathbb{S}_n} |RK_{\rho}(z, r\xi)| d\sigma(\xi) dr \\ = 2n \int_0^1 r^{2n-1} \rho(r) \int_{\mathbb{S}_n} |RK_{\rho}(\xi, rz)| d\sigma(\xi) dr \\ \gtrsim \sum_{d=1}^{\infty} \frac{|z|^d}{\rho_{2n-1+2d}} \int_0^1 r^{2n-1+d} \rho(r) dr \\ = \sum_{d=1}^{\infty} \frac{\rho_{2n-1+d}}{\rho_{2n-1+2d}} |z|^d. \end{aligned}$$

Thus,

$$\begin{aligned} \sup_{z \in \mathbb{B}_n} (1-|z|^2) \int_{\mathbb{B}_n} |RK_{\rho}(z, w)| \rho(w) dv(w) \\ \gtrsim \sup_{z \in \mathbb{B}_n} (1-|z|) \sum_{d=1}^{\infty} \frac{\rho_{d+2n-1}}{\rho_{2d+2n-1}} |z|^d \\ \geq \sup_{N \in \mathbb{N}} \frac{1}{N} \sum_{d=1}^N \frac{\rho_{d+2n-1}}{\rho_{2d+2n-1}} \left(1 - \frac{1}{N}\right)^d \\ \gtrsim \sup_{N \in \mathbb{N}} \frac{1}{N} \sum_{d=1}^N \frac{\rho_{d+2n-1}}{\rho_{2d+2n-1}}. \end{aligned}$$

Since  $P_{\rho}$  is bounded,

$$\sup_{z \in \mathbb{B}_n} (1-|z|^2) \int_{\mathbb{B}_n} |RK_{\rho}(z, w)| \rho(w) dv(w) < \infty.$$

Given  $N \geq 2n$ , we obtain that

$$\begin{aligned} 1 &\gtrsim \frac{1}{4N-2n} \sum_{d=3N-n+1}^{4N-2n} \frac{\rho_{d+2n-1}}{\rho_{2d+2n-1}} \\ &\geq \frac{1}{4N} (N-n) \frac{\rho_{4N}}{\rho_{6N}}. \end{aligned}$$

Hence,

$$\rho_{6N} \gtrsim \rho_{4N}.$$

If  $8N \leq k < 8N+8$ ,  $N \geq 2n+8$ , then

$$\rho_k \leq \rho_{8N} \lesssim \rho_{12N} \lesssim \rho_{18N} \leq \rho_{2k}.$$

On the other hand, since  $\rho$  is positive and integrable on  $[0, 1)$ , it is easy to see that

$$\widehat{\rho}(0) \leq C_0 \widehat{\rho}\left(\frac{1}{2}\right)$$

for some  $C_0 > 0$ .

Hence, by Lemma 2.1 we conclude that  $\rho \in \widehat{\mathcal{D}}$ .  $\square$

From Proposition 3.1 and Proposition 3.2, we obtain the conclusion of Theorem 1.1.

**Remark 3.3.** The method given herein combined with our results in<sup>5</sup> can be used to generalize to the unit ball case the  $L^p$  estimates proved in<sup>7</sup> in the unit disk case. This will be the object of a forthcoming paper.

## Acknowledgement

This study is conducted within the framework of science and technology projects at institutional level of Quy Nhon University under the project code T2021.700.03.

I am grateful to Alexander Borichev and El Hassan Youssfi for helpful remarks.

## REFERENCES

1. O. Constantin, J. A. Peláez. Boundedness of the Bergman projection on  $L^p$  spaces with exponential weights, *Bulletin des Sciences Mathématiques*, **2015**, 139, 245–268.

2. M. Dostanić. Unboundedness of the Bergman projections on  $L^p$  spaces with exponential weights, *Proceedings of the Edinburgh Mathematical Society*, **2004**, 47(1), 111–117.
3. M. Dostanić. Boundedness of the Bergman projections on  $L^p$  spaces with radial weights, *Publications de l'Institut Mathématique*, **2009**, 86, 5–20.
4. P. L. Duren. *Theory of  $H^p$  Spaces*, Academic Press, 1970.
5. V. A. Le. Carleson measures and Toeplitz operators on small Bergman spaces on the ball, *Czechoslovak Mathematical Journal*, **2021**, 71(1), 211–229.
6. J. A. Peláez. Small weighted Bergman spaces, *Proceedings of the summer school in complex and harmonic analysis, and related topics*, Publications of the University of Eastern Finland - Reports and Studies in Forestry and Natural Sciences, Joensuu **2016**, 29–98.
7. J. A. Peláez, J. Rättyä. Two weight inequality for Bergman projection, *Journal de Mathématiques Pures et Appliquées*, **2016**, 105(1), 102–130.
8. J. A. Peláez, J. Rättyä. Bergman projection induced by radial weight, *Advances in Mathematics*, **2021**, 391, 107950.
9. K. Zhu. *Spaces of Holomorphic Functions in the Unit Ball*, Springer-Verlag, New York, 2005.
10. K. Zhu, *Operator Theory in Function Spaces*, 2nd edition, American Mathematical Society, Providence, Rhode Island, 2007.